


# **Ash2 a *Drosophila*: anàlisi funcional i aproximació al complexe**



**Mireia Angulo  
i Parera**

Departament de Genètica  
Facultat de Biologia  
Universitat de Barcelona

**TESI DOCTORAL**

***Ash2 a Drosophila:*  
anàlisi funcional i  
aproximació al complexe**

Mireia Angulo i Parera  
Barcelona, Juny de 2006



# REFERENCES



**A**

**Aasland, R., Gibson, T. J. and Stewart, A. F.** (1995). The PHD finger: implications for chromatin-mediated transcriptional regulation. *Trends Biochem Sci* **20**, 56-9.

**Adams, M. D. Celniker, S. E. Holt, R. A. Evans, C. A. Gocayne, J. D. Amanatides, P. G. Scherer, S. E. Li, P. W. Hoskins, R. A. Galle, R. F. et al.** (2000). The genome sequence of *Drosophila melanogaster*. *Science* **287**, 2185-95.

**Adamson, A. L. and Shearn, A.** (1996). Molecular genetic analysis of *Drosophila* ash2, a member of the trithorax group required for imaginal disc pattern formation. *Genetics* **144**, 621-33.

**Amoros, M.** (2001). Estudio de mutantes del cromosoma III de *Drosophila melanogaster*: el gen ash-2 como regulador de la diferenciación celular. *PhD thesis. Universitat de Barcelona*.

**Amoros, M., Corominas, M., Deak, P. and Serras, F.** (2002). The ash2 gene is involved in *Drosophila* wing development. *Int J Dev Biol* **46**, 321-4.

**Armstrong, J. A., Papoulias, O., Daubresse, G., Sperling, A. S., Lis, J. T., Scott, M. P. and Tamkun, J. W.** (2002). The *Drosophila* BRM complex facilitates global transcription by RNA polymerase II. *Embo J* **21**, 5245-54.

**B**

**Bantignies, F., Goodman, R. H. and Smolik, S. M.** (2000). Functional interaction between the coactivator *Drosophila* CREB-binding protein and ASH1, a member of the trithorax group of chromatin modifiers. *Mol Cell Biol* **20**, 9317-30.

**Barrio, R. and de Celis, J. F.** (2004). Regulation of spalt expression in the *Drosophila* wing blade in response to the Decapentaplegic signaling pathway. *Proc Natl Acad Sci U S A* **101**, 6021-6.

**Basler, K. and Struhl, G.** (1994). Compartment boundaries and the control of *Drosophila* limb pattern by hedgehog protein. *Nature* **368**, 208-14.

**Bate, M. and Arias, A. M.** (1991). The embryonic origin of imaginal discs in *Drosophila*. *Development* **112**, 755-61.

**Beisel, C., Imhof, A., Greene, J., Kremmer, E. and Sauer, F.** (2002). Histone methylation by the *Drosophila* epigenetic transcriptional regulator Ash1. *Nature* **419**, 857-62.

**Beltran, S.** (2006). El transcriptoma d'ash2: dianes i funció. *PhD thesis. Universitat de Barcelona*.

**Beltran, S., Blanco, E., Serras, F., Perez-Villamil, B., Guigo, R., Artavanis-Tsakonas, S. and Corominas, M.** (2003). Transcriptional network controlled by the trithorax-group gene ash2 in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* **100**, 3293-8.

**Bernstein, B. E., Humphrey, E. L., Erlich, R. L., Schneider, R., Bouman, P., Liu, J. S., Kouzarides, T. and Schreiber, S. L.** (2002). Methylation of histone H3 Lys 4 in coding regions of active genes. *Proc Natl Acad Sci U S A* **99**, 8695-700.

**Biehs, B., Sturtevant, M. A. and Bier, E.** (1998). Boundaries in the *Drosophila* wing imaginal disc organize vein-specific genetic programs. *Development* **125**, 4245-57.

**Bienz, M.** (2006). The PHD finger, a nuclear protein-interaction domain. *Trends Biochem Sci* **31**, 35-40.

- Bier, E.** (2000). Drawing lines in the Drosophila wing: initiation of wing vein development. *Curr Opin Genet Dev* **10**, 393-8.
- Brand, A. H. and Perrimon, N.** (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-15.
- Breen, T. R., Chinwalla, V. and Harte, P. J.** (1995). Trithorax is required to maintain engrailed expression in a subset of engrailed-expressing cells. *Mech Dev* **52**, 89-98.
- Brentrup, D., Lerch, H., Jackle, H. and Noll, M.** (2000). Regulation of Drosophila wing vein patterning: net encodes a bHLH protein repressing rhomboid and is repressed by rhomboid-dependent Egfr signalling. *Development* **127**, 4729-41.
- Brizuela, B. J. and Kennison, J. A.** (1997). The Drosophila homeotic gene moira regulates expression of engrailed and HOM genes in imaginal tissues. *Mech Dev* **65**, 209-20.
- Brock, H. W. and Fisher, C. L.** (2005). Maintenance of gene expression patterns. *Dev Dyn* **232**, 633-55.
- Brock, H. W. and van Lohuizen, M.** (2001). The Polycomb group--no longer an exclusive club? *Curr Opin Genet Dev* **11**, 175-81.
- Bulger, M. and Groudine, M.** (1999). Looping versus linking: toward a model for long-distance gene activation. *Genes Dev* **13**, 2465-77.
- Byrd, K. N. and Shearn, A.** (2003). ASH1, a Drosophila trithorax group protein, is required for methylation of lysine 4 residues on histone H3. *Proc Natl Acad Sci U S A* **100**, 11535-40.

## C

- Campbell, G. and Tomlinson, A.** (1999). Transducing the Dpp morphogen gradient in the wing of Drosophila: regulation of Dpp targets by brinker. *Cell* **96**, 553-62.
- Cao, R., Wang, L., Wang, H., Xia, L., Erdjument-Bromage, H., Tempst, P., Jones, R. S. and Zhang, Y.** (2002). Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science* **298**, 1039-43.
- Cavalli, G.** (2002). Chromatin as a eukaryotic template of genetic information. *Curr Opin Cell Biol* **14**, 269-78.
- Cavalli, G. and Paro, R.** (1999). Epigenetic inheritance of active chromatin after removal of the main transactivator. *Science* **286**, 955-8.
- Cheng, M. K. and Shearn, A.** (2004). The direct interaction between ASH2, a Drosophila trithorax group protein, and SKTL, a nuclear phosphatidylinositol 4-phosphate 5-kinase, implies a role for phosphatidylinositol 4,5-bisphosphate in maintaining transcriptionally active chromatin. *Genetics* **167**, 1213-23.
- Chinwalla, V., Jane, E. P. and Harte, P. J.** (1995). The Drosophila trithorax protein binds to specific chromosomal sites and is co-localized with Polycomb at many sites. *Embo J* **14**, 2056-65.
- Cohen, B., Simcox, A. A. and Cohen, S. M.** (1993). Allocation of the thoracic imaginal primordia in the Drosophila embryo. *Development* **117**, 597-608.
- Collins, R. T., Furukawa, T., Tanese, N. and Treisman, J. E.** (1999). Osa associates with the Brahma chromatin remodeling complex and promotes the activation of some target genes. *Embo J* **18**, 7029-40.
- Collins, R. T. and Treisman, J. E.** (2000). Osa-containing Brahma chromatin remodeling complexes are required for the repression of wingless target genes. *Genes Dev* **14**, 3140-52.

**Cook, O., Biehs, B. and Bier, E.** (2004). brinker and optomotor-blind act coordinately to initiate development of the L5 wing vein primordium in *Drosophila*. *Development* **131**, 2113-24.

**Crosby, M. A., Miller, C., Alon, T., Watson, K. L., Verrijzer, C. P., Goldman-Levi, R. and Zak, N. B.** (1999). The trithorax group gene moira encodes a brahma-associated putative chromatin-remodeling factor in *Drosophila melanogaster*. *Mol Cell Biol* **19**, 1159-70.

**Crozatier, M., Glise, B. and Vincent, A.** (2002). Connecting Hh, Dpp and EGF signalling in patterning of the *Drosophila* wing; the pivotal role of collier/knot in the AP organiser. *Development* **129**, 4261-9.

**Czernin, B., Melfi, R., McCabe, D., Seitz, V., Imhof, A. and Pirrotta, V.** (2002). *Drosophila* enhancer of Zeste/ESC complexes have a histone H3 methyltransferase activity that marks chromosomal Polycomb sites. *Cell* **111**, 185-96.

## D

**de Celis, J. F.** (1997). Expression and function of decapentaplegic and thick veins during the differentiation of the veins in the *Drosophila* wing. *Development* **124**, 1007-18.

**De Celis, J. F.** (2003). Pattern formation in the *Drosophila* wing: The development of the veins. *Bioessays* **25**, 443-51.

**de Celis, J. F. and Barrio, R.** (2000). Function of the spalt/spalt-related gene complex in positioning the veins in the *Drosophila* wing. *Mech Dev* **91**, 31-41.

**de Celis, J. F., Barrio, R. and Kafatos, F. C.** (1996). A gene complex acting downstream of dpp in *Drosophila* wing morphogenesis. *Nature* **381**, 421-4.

**de Celis, J. F., Bray, S. and Garcia-Bellido, A.** (1997). Notch signalling regulates veinlet expression and establishes boundaries between veins and interveins in the *Drosophila* wing. *Development* **124**, 1919-28.

**de Celis, J. F. and Garcia-Bellido, A.** (1994). Roles of the Notch gene in *Drosophila* wing morphogenesis. *Mech Dev* **46**, 109-22.

**Deak, P., Omar, M. M., Saunders, R. D., Pal, M., Komonyi, O., Szidonya, J., Maroy, P., Zhang, Y., Ashburner, M., Benos, P. et al.** (1997). P-element insertion alleles of essential genes on the third chromosome of *Drosophila melanogaster*: correlation of physical and cytogenetic maps in chromosomal region 86E-87F. *Genetics* **147**, 1697-722.

**Dejardin, J. and Cavalli, G.** (2004). Chromatin inheritance upon Zeste-mediated Brahma recruitment at a minimal cellular memory module. *Embo J* **23**, 857-68.

**Dejardin, J. and Cavalli, G.** (2005). Epigenetic inheritance of chromatin states mediated by Polycomb and trithorax group proteins in *Drosophila*. *Prog Mol Subcell Biol* **38**, 31-63.

**Diaz-Benjumea, F. J. and Garcia-Bellido, A.** (1990). Behaviour of cells mutant for an EGF receptor homologue of *Drosophila* in genetic mosaics. *Proc Biol Sci* **242**, 36-44.

**Dingwall, A. K., Beek, S. J., McCallum, C. M., Tamkun, J. W., Kalpana, G. V., Goff, S. P. and Scott, M. P.** (1995). The *Drosophila* snr1 and brm proteins are related to yeast SWI/SNF proteins and are components of a large protein complex. *Mol Biol Cell* **6**, 777-91.

**Dominguez, M., Brunner, M., Hafen, E. and Basler, K.** (1996). Sending and receiving the hedgehog signal: control by the *Drosophila* Gli protein Cubitus interruptus. *Science* **272**, 1621-5.

## E

**Elfring, L. K., Daniel, C., Papoulas, O., Deuring, R., Sarte, M., Moseley, S., Beek, S. J., Waldrip, W. R., Daubresse, G., DePace, A. et al.** (1998). Genetic analysis of brahma: the Drosophila homolog of the yeast chromatin remodeling factor SWI2/SNF2. *Genetics* **148**, 251-65.

## F

**Francis, N. J. and Kingston, R. E.** (2001). Mechanisms of transcriptional memory. *Nat Rev Mol Cell Biol* **2**, 409-21.

**Franke, A., DeCamillis, M., Zink, D., Cheng, N., Brock, H. W. and Paro, R.** (1992). Polycomb and polyhomeotic are constituents of a multimeric protein complex in chromatin of Drosophila melanogaster. *Embo J* **11**, 2941-50.

**Fristrom, D., Gotwals, P., Eaton, S., Kornberg, T. B., Sturtevant, M., Bier, E. and Fristrom, J. W.** (1994). Blistered: a gene required for vein/intervein formation in wings of Drosophila. *Development* **120**, 2661-71.

**Funakoshi, Y., Minami, M. and Tabata, T.** (2001). *mtv* shapes the activity gradient of the Dpp morphogen through regulation of thickveins. *Development* **128**, 67-74.

**Furuyama, T., Tie, F. and Harte, P. J.** (2003). Polycomb group proteins ESC and E(Z) are present in multiple distinct complexes that undergo dynamic changes during development. *Genesis* **35**, 114-24.

## G

**Garcia-Bellido, A. and de Celis, J. F.** (1992). Developmental genetics of the venation pattern of Drosophila. *Annu Rev Genet* **26**, 277-304.

**Garcia-Bellido, A., Ripoll, P. and Morata, G.** (1973). Developmental compartmentalisation of the wing disk of Drosophila. *Nat New Biol* **245**, 251-3.

**Gildea, J. J., Lopez, R. and Shearn, A.** (2000). A screen for new trithorax group genes identified little imaginal discs, the Drosophila melanogaster homologue of human retinoblastoma binding protein 2. *Genetics* **156**, 645-63.

**Gomez-Skarmeta, J. L. and Modolell, J.** (1996). araucan and caupolican provide a link between compartment subdivisions and patterning of sensory organs and veins in the Drosophila wing. *Genes Dev* **10**, 2935-45.

**Goo, Y. H., Sohn, Y. C., Kim, D. H., Kim, S. W., Kang, M. J., Jung, D. J., Kwak, E., Barlev, N. A., Berger, S. L., Chow, V. T. et al.** (2003). Activating signal cointegrator 2 belongs to a novel steady-state complex that contains a subset of trithorax group proteins. *Mol Cell Biol* **23**, 140-9.

**Guelman, S., Suganuma, T., Florens, L., Swanson, S. K., Kiesecker, C. L., Kusch, T., Anderson, S., Yates, J. R., 3rd, Washburn, M. P., Abmayr, S. M. et al.** (2006). Host cell factor and an uncharacterized SANT domain protein are stable components of ATAC, a novel dAda2A/dGcn5-containing histone acetyltransferase complex in Drosophila. *Mol Cell Biol* **26**, 871-82.

**Gurdon, J. B. and Bourillot, P. Y.** (2001). Morphogen gradient interpretation. *Nature* **413**, 797-803.

**H**

**Halder, G., Polaczyk, P., Kraus, M. E., Hudson, A., Kim, J., Laughon, A. and Carroll, S.** (1998). The Vestigial and Scalloped proteins act together to directly regulate wing-specific gene expression in Drosophila. *Genes Dev* **12**, 3900-9.

**Hassan, A. H., Prochasson, P., Neely, K. E., Galasinski, S. C., Chandy, M., Carrozza, M. J. and Workman, J. L.** (2002). Function and selectivity of bromodomains in anchoring chromatin-modifying complexes to promoter nucleosomes. *Cell* **111**, 369-79.

**Held, L. I.** (2002). Imaginal Discs: the Genetic and Cellular Logic of Pattern Formation. *Cambridge University Press*.

**Hentze, M. W. and Kulozik, A. E.** (1999). A perfect message: RNA surveillance and nonsense-mediated decay. *Cell* **96**, 307-10.

**Hughes, C. M., Rozenblatt-Rosen, O., Milne, T. A., Copeland, T. D., Levine, S. S., Lee, J. C., Hayes, D. N., Shanmugam, K. S., Bhattacharjee, A., Biondi, C. A. et al.** (2004). Menin associates with a trithorax family histone methyltransferase complex and with the hoxc8 locus. *Mol Cell* **13**, 587-97.

**I**

**Ikegawa, S., Isomura, M., Koshizuka, Y. and Nakamura, Y.** (1999). Cloning and characterization of ASH2L and Ash2l, human and mouse homologs of the Drosophila ash2 gene. *Cytogenet Cell Genet* **84**, 167-72.

**Ingham, P. W.** (1981). Trithorax: a new homeotic mutation of *Drosophila melanogaster*. II. The role of *trx*<sup>+</sup> after embryogenesis. *Wilhelm Roux's Arch Dev Biol* **190**, 365-369.

**Ingham, P. W.** (1983). Differential expression of bithorax complex genes in the absence of the extra sex combs and trithorax genes. *Nature* **306**, 591-593.

**Ingham, P. W.** (1985). A clonal analysis of the requirement for the trithorax gene in the diversification of segments in Drosophila. *J Embryol Exp Morphol* **89**, 349-65.

**Ingham, P. W. and Martinez Arias, A.** (1992). Boundaries and fields in early embryos. *Cell* **68**, 221-35.

**J**

**Jacobs, J. J. and van Lohuizen, M.** (2002). Polycomb repression: from cellular memory to cellular proliferation and cancer. *Biochim Biophys Acta* **1602**, 151-61.

**Jurgens, G.** (1985). A group of genes controlling the spatial expression of the bithorax complex in Drosophila. *Nature* **316**, 153-155.

**K**

**Kal, A. J., Mahmoudi, T., Zak, N. B. and Verrijzer, C. P.** (2000). The Drosophila brahma complex is an essential coactivator for the trithorax group protein zeste. *Genes Dev* **14**, 1058-71.

- Kennison, J. A.** (1995). The Polycomb and trithorax group proteins of Drosophila: trans-regulators of homeotic gene function. *Annu Rev Genet* **29**, 289-303.
- Kennison, J. A. and Tamkun, J. W.** (1988). Dosage-dependent modifiers of polycomb and antennapedia mutations in Drosophila. *Proc Natl Acad Sci U S A* **85**, 8136-40.
- Klymenko, T. and Muller, J.** (2004). The histone methyltransferases Trithorax and Ash1 prevent transcriptional silencing by Polycomb group proteins. *EMBO Rep* **5**, 373-7.
- Krogan, N. J., Dover, J., Khorrami, S., Greenblatt, J. F., Schneider, J., Johnston, M. and Shilatifard, A.** (2002). COMPASS, a histone H3 (Lysine 4) methyltransferase required for telomeric silencing of gene expression. *J Biol Chem* **277**, 10753-5.
- Kurdistani, S. K., Robyr, D., Tavazoie, S. and Grunstein, M.** (2002). Genome-wide binding map of the histone deacetylase Rpd3 in yeast. *Nat Genet* **31**, 248-54.
- Kuzin, B., Tillib, S., Sedkov, Y., Mizrokhi, L. and Mazo, A.** (1994). The Drosophila trithorax gene encodes a chromosomal protein and directly regulates the region-specific homeotic gene fork head. *Genes Dev* **8**, 2478-90.
- Kuzmichev, A., Nishioka, K., Erdjument-Bromage, H., Tempst, P. and Reinberg, D.** (2002). Histone methyltransferase activity associated with a human multiprotein complex containing the Enhancer of Zeste protein. *Genes Dev* **16**, 2893-905.

## L

- LaJeunesse, D. and Shearn, A.** (1995). Trans-regulation of thoracic homeotic selector genes of the Antennapedia and bithorax complexes by the trithorax group genes: absent, small, and homeotic discs 1 and 2. *Mech Dev* **53**, 123-39.
- Lee, J. H. and Skalnik, D. G.** (2005). CpG-binding protein (CXXC finger protein 1) is a component of the mammalian Set1 histone H3-Lys4 methyltransferase complex, the analogue of the yeast Set1/COMPASS complex. *J Biol Chem* **280**, 41725-31.
- Lewis, E. B.** (1998). The bithorax complex: the first fifty years. *Int J Dev Biol* **42**, 403-15.
- Lunde, K., Biehs, B., Nauber, U. and Bier, E.** (1998). The knirps and knirps-related genes organize development of the second wing vein in Drosophila. *Development* **125**, 4145-54.
- Lunde, K., Trimble, J. L., Guichard, A., Guss, K. A., Nauber, U. and Bier, E.** (2003). Activation of the knirps locus links patterning to morphogenesis of the second wing vein in Drosophila. *Development* **130**, 235-48.

## M

- Mahajan, S. S., Johnson, K. M. and Wilson, A. C.** (2003). Molecular cloning of Drosophila HCF reveals proteolytic processing and self-association of the encoded protein. *J Cell Physiol* **194**, 117-26.
- Marenda, D. R., Zraly, C. B., Feng, Y., Egan, S. and Dingwall, A. K.** (2003). The Drosophila SNR1 (SNF5/INI1) subunit directs essential developmental functions of the Brahma chromatin remodeling complex. *Mol Cell Biol* **23**, 289-305.

- Martin-Blanco, E., Roch, F., Noll, E., Baonza, A., Duffy, J. B. and Perrimon, N.** (1999). A temporal switch in DER signaling controls the specification and differentiation of veins and interveins in the Drosophila wing. *Development* **126**, 5739-47.
- Matakatsu, H., Tadokoro, R., Gamo, S. and Hayashi, S.** (1999). Repression of the wing vein development in Drosophila by the nuclear matrix protein plexus. *Development* **126**, 5207-16.
- McKeon, J., Slade, E., Sinclair, D. A., Cheng, N., Couling, M. and Brock, H. W.** (1994). Mutations in some Polycomb group genes of Drosophila interfere with regulation of segmentation genes. *Mol Gen Genet* **244**, 474-83.
- Miller, T., Krogan, N. J., Dover, J., Erdjument-Bromage, H., Tempst, P., Johnston, M., Greenblatt, J. F. and Shilatifard, A.** (2001). COMPASS: a complex of proteins associated with a trithorax-related SET domain protein. *Proc Natl Acad Sci U S A* **98**, 12902-7.
- Minami, M., Kinoshita, N., Kamoshida, Y., Tanimoto, H. and Tabata, T.** (1999). brinker is a target of Dpp in Drosophila that negatively regulates Dpp-dependent genes. *Nature* **398**, 242-6.
- Mohler, J., Seecoomar, M., Agarwal, S., Bier, E. and Hsai, J.** (2000). Activation of knot (kn) specifies the 3-4 intervein region in the Drosophila wing. *Development* **127**, 55-63.
- Mohrmann, L., Langenberg, K., Krijgsveld, J., Kal, A. J., Heck, A. J. and Verrijzer, C. P.** (2004). Differential targeting of two distinct SWI/SNF-related Drosophila chromatin-remodeling complexes. *Mol Cell Biol* **24**, 3077-88.
- Montagne, J., Groppe, J., Guillemin, K., Krasnow, M. A., Gehring, W. J. and Affolter, M.** (1996). The Drosophila Serum Response Factor gene is required for the formation of intervein tissue of the wing and is allelic to blistered. *Development* **122**, 2589-97.
- Morata, G. and Ripoll, P.** (1975). Minutes: mutants of drosophila autonomously affecting cell division rate. *Dev Biol* **42**, 211-21.
- Muller, B., Hartmann, B., Pyrowolakis, G., Affolter, M. and Basler, K.** (2003). Conversion of an extracellular Dpp/BMP morphogen gradient into an inverse transcriptional gradient. *Cell* **113**, 221-33.
- Muller, J., Hart, C. M., Francis, N. J., Vargas, M. L., Sengupta, A., Wild, B., Miller, E. L., O'Connor, M. B., Kingston, R. E. and Simon, J. A.** (2002). Histone methyltransferase activity of a Drosophila Polycomb group repressor complex. *Cell* **111**, 197-208.

## N

- Nagy, P. L., Griesenbeck, J., Kornberg, R. D. and Cleary, M. L.** (2002). A trithorax-group complex purified from *Saccharomyces cerevisiae* is required for methylation of histone H3. *Proc Natl Acad Sci U S A* **99**, 90-4.
- Nakamura, T., Mori, T., Tada, S., Krajewski, W., Rozovskaia, T., Wassell, R., Dubois, G., Mazo, A., Croce, C. M. and Canaani, E.** (2002). ALL-1 is a histone methyltransferase that assembles a supercomplex of proteins involved in transcriptional regulation. *Mol Cell* **10**, 1119-28.
- Ng, J., Hart, C. M., Morgan, K. and Simon, J. A.** (2000). A Drosophila ESC-E(Z) protein complex is distinct from other polycomb group complexes and contains covalently modified ESC. *Mol Cell Biol* **20**, 3069-78.

**Nussbaumer, U., Halder, G., Groppe, J., Affolter, M. and Montagne, J.** (2000). Expression of the blistered/DSRF gene is controlled by different morphogens during Drosophila trachea and wing development. *Mech Dev* **96**, 27-36.

## O

**Orlando, V.** (2003). Polycomb, epigenomes, and control of cell identity. *Cell* **112**, 599-606.

**Orlando, V., Jane, E. P., Chinwalla, V., Harte, P. J. and Paro, R.** (1998). Binding of trithorax and Polycomb proteins to the bithorax complex: dynamic changes during early Drosophila embryogenesis. *Embo J* **17**, 5141-50.

**Orphanides, G., LeRoy, G., Chang, C. H., Luse, D. S. and Reinberg, D.** (1998). FACT, a factor that facilitates transcript elongation through nucleosomes. *Cell* **92**, 105-16.

**Otte, A. P. and Kwaks, T. H.** (2003). Gene repression by Polycomb group protein complexes: a distinct complex for every occasion? *Curr Opin Genet Dev* **13**, 448-54.

## P

**Papoulias, O., Beek, S. J., Moseley, S. L., McCallum, C. M., Sarte, M., Shearn, A. and Tamkun, J. W.** (1998). The Drosophila trithorax group proteins BRM, ASH1 and ASH2 are subunits of distinct protein complexes. *Development* **125**, 3955-66.

**Paumard-Rigal, S., Zider, A., Vaudin, P. and Silber, J.** (1998). Specific interactions between vestigial and scalloped are required to promote wing tissue proliferation in Drosophila melanogaster. *Dev Genes Evol* **208**, 440-6.

**Pelegrí, F. and Lehmann, R.** (1994). A role of polycomb group genes in the regulation of gap gene expression in Drosophila. *Genetics* **136**, 1341-53.

**Petruk, S., Sedkov, Y., Smith, S., Tillib, S., Kraevski, V., Nakamura, T., Canaani, E., Croce, C. M. and Mazo, A.** (2001). Trithorax and dCBP acting in a complex to maintain expression of a homeotic gene. *Science* **294**, 1331-4.

**Pile, L. A., Spellman, P. T., Katzenberger, R. J. and Wassarman, D. A.** (2003). The SIN3 deacetylase complex represses genes encoding mitochondrial proteins: implications for the regulation of energy metabolism. *J Biol Chem* **278**, 37840-8.

**Pile, L. A. and Wassarman, D. A.** (2000). Chromosomal localization links the SIN3-RPD3 complex to the regulation of chromatin condensation, histone acetylation and gene expression. *Embo J* **19**, 6131-40.

**Ponting, C., Schultz, J. and Bork, P.** (1997). SPRY domains in ryanodine receptors (Ca(2+)-release channels). *Trends Biochem Sci* **22**, 193-4.

**Poux, S., Horard, B., Sigrist, C. J. and Pirrotta, V.** (2002). The Drosophila trithorax protein is a coactivator required to prevent re-establishment of polycomb silencing. *Development* **129**, 2483-93.

**Poux, S., McCabe, D. and Pirrotta, V.** (2001a). Recruitment of components of Polycomb Group chromatin complexes in Drosophila. *Development* **128**, 75-85.

**Poux, S., Melfi, R. and Pirrotta, V.** (2001b). Establishment of Polycomb silencing requires a transient interaction between PC and ESC. *Genes Dev* **15**, 2509-14.

**R**

**Ragvin, A., Valvatne, H., Erdal, S., Arskog, V., Tufteland, K. R., Breen, K., AM, O. Y., Eberharter, A., Gibson, T. J., Becker, P. B. et al.** (2004). Nucleosome binding by the bromodomain and PHD finger of the transcriptional cofactor p300. *J Mol Biol* **337**, 773-88.

**Rank, G., Prestel, M. and Paro, R.** (2002). Transcription through intergenic chromosomal memory elements of the Drosophila bithorax complex correlates with an epigenetic switch. *Mol Cell Biol* **22**, 8026-34.

**Rastelli, L., Chan, C. S. and Pirrotta, V.** (1993). Related chromosome binding sites for zeste, suppressors of zeste and Polycomb group proteins in Drosophila and their dependence on Enhancer of zeste function. *Embo J* **12**, 1513-22.

**Ringrose, L. and Paro, R.** (2004). Epigenetic regulation of cellular memory by the Polycomb and Trithorax group proteins. *Annu Rev Genet* **38**, 413-43.

**Ringrose, L., Rehmsmeier, M., Dura, J. M. and Paro, R.** (2003). Genome-wide prediction of Polycomb/Trithorax response elements in *Drosophila melanogaster*. *Dev Cell* **5**, 759-71.

**Roch, F., Baonza, A., Martin-Blanco, E. and Garcia-Bellido, A.** (1998). Genetic interactions and cell behaviour in blistered mutants during proliferation and differentiation of the *Drosophila* wing. *Development* **125**, 1823-32.

**Roguev, A., Schaft, D., Shevchenko, A., Aasland, R., Shevchenko, A. and Stewart, A. F.** (2003). High conservation of the Set1/Rad6 axis of histone 3 lysine 4 methylation in budding and fission yeasts. *J Biol Chem* **278**, 8487-93.

**Roguev, A., Schaft, D., Shevchenko, A., Pijnappel, W. W., Wilm, M., Aasland, R. and Stewart, A. F.** (2001). The *Saccharomyces cerevisiae* Set1 complex includes an Ash2 homologue and methylates histone 3 lysine 4. *Embo J* **20**, 7137-48.

**Rozovskaia, T., Tillib, S., Smith, S., Sedkov, Y., Rozenblatt-Rosen, O., Petruk, S., Yano, T., Nakamura, T., Ben-Simchon, L., Gildea, J. et al.** (1999). Trithorax and ASH1 interact directly and associate with the trithorax group-responsive bxd region of the Ultrabithorax promoter. *Mol Cell Biol* **19**, 6441-7.

**Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J., Webster, J., Gubb, D., Gunton, N., Johnson, G. et al.** (2004). The DrosDel collection: a set of P-element insertions for generating custom chromosomal aberrations in *Drosophila melanogaster*. *Genetics* **167**, 797-813.

**S**

**Saget, O., Forquignon, F., Santamaria, P. and Randsholt, N. B.** (1998). Needs and targets for the multi sex combs gene product in *Drosophila melanogaster*. *Genetics* **149**, 1823-38.

**Sanchez-Elsner, T., Gou, D., Kremmer, E. and Sauer, F.** (2006). Noncoding RNAs of trithorax response elements recruit *Drosophila* Ash1 to Ultrabithorax. *Science* **311**, 1118-23.

**Santos-Rosa, H., Schneider, R., Bannister, A. J., Sherriff, J., Bernstein, B. E., Emre, N. C., Schreiber, S. L., Mellor, J. and Kouzarides, T.** (2002). Active genes are tri-methylated at K4 of histone H3. *Nature* **419**, 407-11.

**Satijn, D. P. and Otte, A. P.** (1999). Polycomb group protein complexes: do different complexes regulate distinct target genes? *Biochim Biophys Acta* **1447**, 1-16.

- Saurin, A. J., Shao, Z., Erdjument-Bromage, H., Tempst, P. and Kingston, R. E.** (2001). A Drosophila Polycomb group complex includes Zeste and dTAFII proteins. *Nature* **412**, 655-60.
- Schmitt, S., Prestel, M. and Paro, R.** (2005). Intergenic transcription through a polycomb group response element counteracts silencing. *Genes Dev* **19**, 697-708.
- Schneider, J., Wood, A., Lee, J. S., Schuster, R., Dueker, J., Maguire, C., Swanson, S. K., Florens, L., Washburn, M. P. and Shilatifard, A.** (2005). Molecular regulation of histone H3 trimethylation by COMPASS and the regulation of gene expression. *Mol Cell* **19**, 849-56.
- Schnepf, B., Grumblig, G., Donaldson, T. and Simcox, A.** (1996). Vein is a novel component in the Drosophila epidermal growth factor receptor pathway with similarity to the neuregulins. *Genes Dev* **10**, 2302-13.
- Shao, Z., Raible, F., Mollaaghbabab, R., Guyon, J. R., Wu, C. T., Bender, W. and Kingston, R. E.** (1999). Stabilization of chromatin structure by PRC1, a Polycomb complex. *Cell* **98**, 37-46.
- Shearn, A.** (1989). The ash-1, ash-2 and trithorax genes of Drosophila melanogaster are functionally related. *Genetics* **121**, 517-25.
- Shearn, A., Hersperger, E. and Hersperger, E.** (1987). Genetic studies of mutations at two loci of Drosophila melanogaster which cause a wide variety of homeotic transformations. *Roux's Arch Dev Biol* **196**, 231-42.
- Shearn, A., Rice, T., Garen, A. and Gehring, W.** (1971). Imaginal disc abnormalities in lethal mutants of Drosophila. *Proc Natl Acad Sci U S A* **68**, 2594-8.
- Sierra, J., Yoshida, T., Joazeiro, C. A. and Jones, K. A.** (2006). The APC tumor suppressor counteracts beta-catenin activation and H3K4 methylation at Wnt target genes. *Genes Dev* **20**, 586-600.
- Silverstein, R. A. and Ekwall, K.** (2005). Sin3: a flexible regulator of global gene expression and genome stability. *Curr Genet* **47**, 1-17.
- Simcox, A. A., Grumblig, G., Schnepf, B., Bennington-Mathias, C., Hersperger, E. and Shearn, A.** (1996). Molecular, phenotypic, and expression analysis of vein, a gene required for growth of the Drosophila wing disc. *Dev Biol* **177**, 475-89.
- Simon, J.** (1995). Locking in stable states of gene expression: transcriptional control during Drosophila development. *Curr Opin Cell Biol* **7**, 376-85.
- Smith, S. T., Petruk, S., Sedkov, Y., Cho, E., Tillib, S., Canaani, E. and Mazo, A.** (2004). Modulation of heat shock gene expression by the TAC1 chromatin-modifying complex. *Nat Cell Biol* **6**, 162-7.
- Sotillos, S. and De Celis, J. F.** (2005). Interactions between the Notch, EGFR, and decapentaplegic signaling pathways regulate vein differentiation during Drosophila pupal wing development. *Dev Dyn* **232**, 738-52.
- Strahl, B. D. and Allis, C. D.** (2000). The language of covalent histone modifications. *Nature* **403**, 41-5.
- Struuli, P. and Arnosti, D. N.** (2005). Functional interaction between the Drosophila knirps short range transcriptional repressor and RPD3 histone deacetylase. *J Biol Chem* **280**, 40757-65.
- Struhl, G. and Akam, M.** (1985). Altered distributions of Ultrabithorax transcripts in extra sex combs mutant embryos of Drosophila. *Embo J* **4**, 3259-64.
- Sturtevant, M. A., Biehs, B., Marin, E. and Bier, E.** (1997). The spalt gene links the A/P compartment boundary to a linear adult structure in the Drosophila wing. *Development* **124**, 21-32.
- Sturtevant, M. A. and Bier, E.** (1995). Analysis of the genetic hierarchy guiding wing vein development in Drosophila. *Development* **121**, 785-801.

**Sturtevant, M. A., Roark, M. and Bier, E.** (1993). The Drosophila rhomboid gene mediates the localized formation of wing veins and interacts genetically with components of the EGF-R signaling pathway. *Genes Dev* **7**, 961-73.

## T

**Tabata, T. and Kornberg, T. B.** (1994). Hedgehog is a signaling protein with a key role in patterning Drosophila imaginal discs. *Cell* **76**, 89-102.

**Tanimoto, H., Itoh, S., ten Dijke, P. and Tabata, T.** (2000). Hedgehog creates a gradient of DPP activity in Drosophila wing imaginal discs. *Mol Cell* **5**, 59-71.

**Teleman, A. A. and Cohen, S. M.** (2000). Dpp gradient formation in the Drosophila wing imaginal disc. *Cell* **103**, 971-80.

**Tie, F., Furuyama, T., Prasad-Sinha, J., Jane, E. and Harte, P. J.** (2001). The Drosophila Polycomb Group proteins ESC and E(Z) are present in a complex containing the histone-binding protein p55 and the histone deacetylase RPD3. *Development* **128**, 275-86.

**Tie, F., Prasad-Sinha, J., Birve, A., Rasmuson-Lestander, A. and Harte, P. J.** (2003). A 1-megadalton ESC/E(Z) complex from Drosophila that contains polycomblike and RPD3. *Mol Cell Biol* **23**, 3352-62.

**Tillib, S., Petruk, S., Sedkov, Y., Kuzin, A., Fujioka, M., Goto, T. and Mazo, A.** (1999). Trithorax- and Polycomb-group response elements within an Ultrabithorax transcription maintenance unit consist of closely situated but separable sequences. *Mol Cell Biol* **19**, 5189-202.

**Tolhuis, B., Muijrers, I., de Wit, E., Teunissen, H., Talhout, W., van Steensel, B. and van Lohuizen, M.** (2006). Genome-wide profiling of PRC1 and PRC2 Polycomb chromatin binding in Drosophila melanogaster. *Nat Genet*.

**Treisman, J. E., Luk, A., Rubin, G. M. and Heberlein, U.** (1997). eyelid antagonizes wingless signaling during Drosophila development and has homology to the Bright family of DNA-binding proteins. *Genes Dev* **11**, 1949-62.

**Tripoulas, N., LaJeunesse, D., Gildea, J. and Shearn, A.** (1996). The Drosophila ash1 gene product, which is localized at specific sites on polytene chromosomes, contains a SET domain and a PHD finger. *Genetics* **143**, 913-28.

**Turner, F. R. and Mahowald, A. P.** (1979). Scanning electron microscopy of Drosophila melanogaster embryogenesis. III. Formation of the head and caudal segments. *Dev Biol* **68**, 96-109.

## V

**van Lohuizen, M.** (1999). The trithorax-group and polycomb-group chromatin modifiers: implications for disease. *Curr Opin Genet Dev* **9**, 355-61.

**Vermaak, D., Ahmad, K. and Henikoff, S.** (2003). Maintenance of chromatin states: an open-and-shut case. *Curr Opin Cell Biol* **15**, 266-74.

**Vervoort, M., Crozatier, M., Valle, D. and Vincent, A.** (1999). The COE transcription factor Collier is a mediator of short-range Hedgehog-induced patterning of the Drosophila wing. *Curr Biol* **9**, 632-9.

**Vidal, M. and Gaber, R. F.** (1991). RPD3 encodes a second factor required to achieve maximum positive and negative transcriptional states in *Saccharomyces cerevisiae*. *Mol Cell Biol* **11**, 6317-27.

**Vidal, M., Strich, R., Esposito, R. E. and Gaber, R. F.** (1991). RPD1 (SIN3/UME4) is required for maximal activation and repression of diverse yeast genes. *Mol Cell Biol* **11**, 6306-16.

## W

**Wang, H., Wang, L., Erdjument-Bromage, H., Vidal, M., Tempst, P., Jones, R. S. and Zhang, Y.** (2004a). Role of histone H2A ubiquitination in Polycomb silencing. *Nature* **431**, 873-8.

**Wang, J., Zhou, Y., Yin, B., Du, G., Huang, X., Li, G., Shen, Y., Yuan, J. and Qiang, B.** (2001). ASH2L: alternative splicing and downregulation during induced megakaryocytic differentiation of multipotential leukemia cell lines. *J Mol Med* **79**, 399-405.

**Wang, L., Brown, J. L., Cao, R., Zhang, Y., Kassis, J. A. and Jones, R. S.** (2004b). Hierarchical recruitment of polycomb group silencing complexes. *Mol Cell* **14**, 637-46.

**Williams, J. A., Paddock, S. W. and Carroll, S. B.** (1993). Pattern formation in a secondary field: a hierarchy of regulatory genes subdivides the developing Drosophila wing disc into discrete subregions. *Development* **117**, 571-84.

**Wilson, A. C., Peterson, M. G. and Herr, W.** (1995). The HCF repeat is an unusual proteolytic cleavage signal. *Genes Dev* **9**, 2445-58.

**Wysocka, J., Myers, M. P., Laherty, C. D., Eisenman, R. N. and Herr, W.** (2003). Human Sin3 deacetylase and trithorax-related Set1/Ash2 histone H3-K4 methyltransferase are tethered together selectively by the cell-proliferation factor HCF-1. *Genes Dev* **17**, 896-911.

## X

**Xu, T. and Rubin, G. M.** (1993). Analysis of genetic mosaics in developing and adult Drosophila tissues. *Development* **117**, 1223-37.

## Y

**Yokoyama, A., Wang, Z., Wysocka, J., Sanyal, M., Aufiero, D. J., Kitabayashi, I., Herr, W. and Cleary, M. L.** (2004). Leukemia proto-oncoprotein MLL forms a SET1-like histone methyltransferase complex with menin to regulate Hox gene expression. *Mol Cell Biol* **24**, 5639-49.

**Yu, K., Sturtevant, M. A., Biehs, B., Francois, V., Padgett, R. W., Blackman, R. K. and Bier, E.** (1996). The Drosophila decapentaplegic and short gastrulation genes function antagonistically during adult wing vein development. *Development* **122**, 4033-44.

AGRAiMENTS



En principi no tenia pensat fer uns agraïments massa extensos, i no pel fet que no tingui coses a agrair sinó per tot el contrari. Tinc tantes coses per agrair que em fa por deixar-me gent per anomenar, per això aquests agraïments van dedicats a TOTHOM que sempre ha estat allà quan ho he necessitat. Primer de tot, he d'agrair-li a la meva família..... TOT, però especialment el suport que m'han donat sempre, i també li he d'agrair al Sergi que aquesta Tesi s'hagi dipositat en el seu moment, i a més, que s'hagi dipositat amb els espais correctes i les lletres en cursives, o no, que toca. Almenys això espero! De veritat, moltes gràcies.

A qui no puc deixar d'anomenar són als meus "jefes" i als meus companys de lab. A la Montse li agraeixo que sigui una persona molt pràctica i sàpiga dir-te: això no cal, o això sí, i al Florenci li agraeixo que sempre m'hagi escoltat en els meus moments de no saber ni el per què, ni el com ni el quan. Seguidament, als meus companys de lab. Des de que vaig entrar he d'agrair al Manel, el meu guia espiritual, a la Marta, la meva companya de lab, de festa i de viatges, a la Cris, per fer-me riure amb les seves sortides, a l'Adrià, no només per tot el que m'ha ajudat amb la feina sinó per haver compartit bons moments, a la Isabel, per no queixar-se de la meva invasió de papers cap a la seva taula i a l'Albert, la nova incorporació que promete.

Dels pisos de dalt m'agradaria agrair a tots aquells, que són molts, que m'han ajudat amb els temes moleculars, especialment a la Gemma i a la Neus, i del meu pis i del Depart en general, crec que li dono les gràcies a tothom per ser com són.



