

Chapter 9

Bibliography

Abrahamsen B, Hansen TB, Hosberg IM (1996): Impact of hemodialysis on dual X-ray absorptiometry, bioelectrical impedance measurements, and anthropometry. *Am J Clin Nutr*, **63**:80-86.

Ainsworth BE, Stolarczyk LM and Heyward VH (1997): Predictive accuracy of bioimpedance in estimating fat-free mass of African-American women. *Med Sci Sports Exerc*, **29**:781.

Altman DG, Machin D, Bryant TN, Gardner MJ (2000): *Statistics with confidence* (2nd ed). Bristol, British Medical Journal Books.

Al-Hatib F (1998): Patient-Instrument connection errors in bioelectrical impedance measurement. *Physiol Meas*, **19**:285-296.

Arkouche W, Fouque D, Pachiaudi C (1997): Total body water and body composition in chronic peritoneal dialysis patients. *J Am Soc Nephrol*, **8**: 1906-1914.

Armitage P, Berry G, Matthews JNS (2002): *Statistical Methods in medical research*. Blackwell Science, 4th ed, London, England.

Armstrong LE, Kenefick RW, Castellani JW, Riebe D, Kavouras SA, Kuznicki JT, Maresh CM (1997): Bioimpedance spectroscopy technique: Intra, extra and total body water. *Med Sci Sports Exerc*, **29**(12): 1657-1663.

Bandini LG, Vu DM, Must A, Dietz WH (1997): Body fatness and bioelectrical impedance in non-obese pre-menarcheal girls: comparison to anthropometry and evaluation of predictive equations. *Eur J Clin Nutr*, **51**:673-677.

Battistini N, Bedogni G, Mariazi E, Severi S, Andreoli A, de Lorenzo A (1996): Relationship between bioelectric impedance, body muscularity and body adiposity in young children. *Riv Ital Pediatr*, **22**:47-53.

Baumgartner RN (1996): Electrical impedance and total body electrical conductivity. In: Roche AF, Heymsfield SB, Lohman TG, eds. *Human body composition*. Champaign, IL: Human Kinetics, 79-107.

Baumgartner RN, Chumlea WC, Roche AF (1989): Estimation of body composition from bioelectric impedance of body segments. *Am J Clin Nutr*, **50**:221-226

Belalcazar A, Patterson R (2004): Improved lung edema monitoring with coronary vein pacing leads: a simulation study. *Physiol Meas*, **25**:475-487.

Bioelectrical impedance analysis in body composition measurement (1994): *National Institutes of Health Technology Assessment Conference Statement*. December 12-14, 1 – 35.

Bishay ST (2000): Numerical methods for the calculation of the Cole-Cole parameters. *Egypt J Sol* **23**(2):179-188.

Bland JM, Altman DG (1986): Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*, **1**:307-310.

- Bland JM, Altman DG (1999): Measuring agreement in method comparison studies. *Statistical Methods in Medical Research*, **8**:135-160.
- Bolton MP, Ward LC, Khan A, Campbell I, Nightingale P, Dewit O, Elia M (1998): Sources of error in bioimpedance spectroscopy. *Physiol Meas*, **19**(2): 235-245.
- Box George EP, Hunter WG, Hunter JS (2001) : Estadística para investigadores. Editorial Reverté, SA, 2001
- Borawski J, Pawlak K, Mysliwiec M (2002): Inflammatory markers and platelet aggregation test as predictor of hemoglobin and endogenous erythropoietin levels in hemodialysis patients. *Nephron*, **91**: 671-681.
- Bradbury MG, Smye SW, Brocklebank JT (2001): Measurement on intercompartmental fluid shifts during hemodialysis in children. *Physiol. Meas*, **22**:351-363.
- Bracco D, Thiébaud D, Chioléro RL (1996): Segmental body composition assessed by bioelectrical impedance analysis and DEXA in humans. *J. Appl. Physiol*, **81**(6): 2580-2587.
- Brodie D, Moscrip V, Hutcheon R (1998): Body composition measurement: a review of hydrodensitometry, anthropometry and impedance methods. *Nutrition*, **14**(3): 296-310
- Brown BH, Barber DC, Morice AH (1994): Cardiac and respiratory related electrical impedance changes in the human thorax. *IEEE Transactions on Biomedical Engineering*, **41**(8): 729-733.
- Bulygin VP, Chepaikin AG (2003): Determination of sensitivity and specificity parameters of interpretation rules in medical device testing. *Biomedical Engineering*, **37**(6):309-314.
- Cai Y, Jenstrup M, Ide K, Perko M, Secher NH (2000): Influence of temperature on the distribution of blood in humans as assessed by electrical impedance. *Eur J Appl Physiol*, **81**:443-448
- Cai Y, Zimmerman A, Ladefoged S (2002): Can hemodialysis-induced hypotension be predicted ? *Nephron*, **92** :582-588.
- Carela MJ, Rodgers CD, Anderson D, Gossain VV (1997): Serial measurements of body composition in obese subjects during a very low energy diet (VLED) comparing bioelectrical impedance with hydrodensitometry. *Obes Res*, **5**(3):250-256.
- Carter M, Morris AT, Zhu F, Zaluska W and Levin NW (2005): Effect of body mass index (BMI) on estimation of extracellular volume (ECV) in hemodialysis (HD) patients using segmental and whole body bioimpedance analysis. *Physiol Meas*, **26**:S93-S99.
- Catalano PM, Wong WW, Drago NM, Amini SB (1995): Estimating body composition in late gestation: a new hydration constant for body density and total body water. *Am J Physiol*, **31**(268): E153-E158
- Cole KS (1940): Permeability and impermeability of cell membranes for ions. *Cold Spring Harbor Sympos Quant Biol*, **8**, 110-122.
- Cole KS (1932): Electrical phase angle of cell membranes. *J Gen Physiol*, **15**, 641-649.
- Cole KS (1928-b): Electrical impedance of suspensions of arbacia eggs. *J Gen Physiol*, **12**, 37-54.
- Cole KS (1928-a): Electrical impedance of suspensions of spheres. *J Gen Physiol*, **12**, 29-36.
- Cole KS (1972): Membranes, ions and impulses: a chapter of classical biophysics. Berkeley, CA: University of California Press.

- Cooper BA, Aslani A, Ryan M (2000): Comparing different methods of assessing body composition in end-stage renal failure. *Kidney Int*, **58**:408-416.
- Cox-Reijven PLM, Van Kreel B, Soeters PB (2002): Bio-electrical impedance spectroscopy: Alternatives for the conventional hand-to-foot measurements. *Clinical Nutrition*, **21** (2): 127-133.
- Cooter G, Moshkovitz Y, Kaluski E, Cohen AJ (2004): Accurate, non-invasive continuous monitoring of cardiac output by whole-body electrical bioimpedance. *CHEST*, **125**:1431-1440.
- Cornish BH, Jacobs A, Thomas BJ, Ward LC (1999): Optimizing electrode sites for segmental bioimpedance measurements. *Physiol Meas*, **20**:241-250.
- Cornish BH, Ward LC, Thomas BJ, Jebb SA, Elia M (1996): Evaluation of multiple frequency bioelectrical impedance and Cole-Cole analysis for the assessment of body water volumes in healthy humans. *Eur J Clin Nutr*, **50**:159-164.
- Cornish BH, Ward LC, Thomas BJ (1994): Alteration of the extracellular and total body water volumes measured by multiple frequency bioelectrical impedance analysis (MFBIA). *Nutr Research*, **14**(5):717-727.
- Cornish BH, Thomas BJ, Ward LC (1993): Improved prediction of extracellular and total body water using impedance loci generated by multiple frequency bioelectrical impedance analysis. *Phys Med Biol*, **38**:337.
- Critchley LAH, Calcroft RM, Tan PYH (2000): The effect of lung injury and excessive lung fluid, on impedance cardiac output measurements, in the critically ill. *Intensive. Care. Med*, **26**:679-685.
- Critchley LAH, Critchley JAJH (1998): Case Report. Lung fluid and impedance cardiography. *Anesthesi*, **53**: 369-381.
- Critchley LAH, Critchley JAJH (1997): Wet lungs and impedance cardiography. *Anesthesia and Int Care*, **25**:314-315.
- Chanchairujira T, Metha RL (2001): Assessing fluid change in hemodialysis: Whole body versus sum of segmental bioimpedance spectroscopy. *Kidney Int*, **60**: 2337-2342.
- Charach G, Rabinovich P, Grosskopf I (2001): Transthoracic monitoring of the impedance of the right lung in patients with cardiogenic pulmonary edema. *Crit Care Med*, **29**(6): 1137-1144.
- Chen YC, Chen HH, Yeh JC, Chen SY (2002): Adjusting dry weight by extracellular volume and body composition in hemodialysis patients. *Nephron*, **92**(1):91-6.
- Chertow GM, Johansen KL, Lew NL (2000): Vintage, nutritional status, and survival in hemodialysis patients: *Kidney Inter*, **57**:1176-1181.
- Chertow GM, Lazarus JM, Lew NL (1997): Development of a population-specific regression equation to estimate total body water in hemodialysis patients. *Kidney Int*, **51**:1578-1582.
- Chertow GM, Johansen KL, Lew NL (1997): Bioimpedance norms for the hemodialysis population: *Kidney Inter*. **52**:1617-1621.
- Chertow GM, Jacobs DO, Lazarus JM (1997): Phase angle predicts survival in hemodialysis patients. *J. Renal Nutr*, **7**:204-207.
- Chertow GM, Lowrie EG, Lew NL (1996): Bioelectrical impedance analysis predicts survival in hemodialysis patients. *J Am Soc Nephrol*, **7**:1442.
- Chertow GM, Lowrie EG, Wilmore DW (1995): Nutritional assessment with bioelectrical impedance analysis in maintenance hemodialysis patients. *J Am Soc Neph*, **6**:75-81.

Chumlea WC, Guo SS, Zeller CM (2001): Total body water reference values and prediction equations for adults. *Kidney Int*, **59**:2250-2258.

Chumlea WC, Guo SS, Zeller CM, Reo NV, Siervogel RM (1999): Total body water data for white adults 18 to 64 years of age: The Fels longitudinal study. *Kidney Int*, **56**:244-252.

Chumlea WC, Guo SS, Siervogel RM (1998): Phase angle spectrum analysis and body water. *Appl Radiat Isot*, **49**(5/6):489-491

Chumlea WC, Guo SS (1997): Bioelectrical Impedance: A history, research issues and recent consensus. In: *Emerging technologies for nutrition research*, National Academy Press, Washington, DC, 169-192.

Chumlea WC, Guo SS, Cockram DB, Siervogel RM (1996): Mechanical and physiological modifiers and bioelectrical impedance spectrum determinants of body composition. *Am J Clin Nutr*, **64**(suppl):413S-422S.

Chumlea WC, Guo SS (1994): Bioelectrical impedance and body composition: present status and future directions. *Nutr Reviews*, **52**(4):123-131.

Chumlea WC, Baumgartner RN, Roche AF (1988): Specific resistivity used to estimate fat-free mass from segmental body measures of bioelectrical impedance. *Am J Clin Nutr*, **48**:7-15

Daugirdas JT, Blake PG, Ing TS (2001): *Handbook of dialysis* (3rd ed). Philadelphia, Lippincott Williams & Wilkins.

De Lorenzo A, di Campli C, Andreoli A, Sasso GF, Bonamico M, Gasbarrini A (1999): Assessment of body composition by bioelectrical impedance in adolescent patients with Celiac disease. *Am J Gastroenter*, **94**(10):2951-2955.

De Lorenzo A, Sorge RP, Candeloro N, di Campli C, Sesti G, Lauro R (1999): New insights into body composition assessment in obese women. *Can J Physiol Pharmacol*, **77**:17-21.

De Lorenzo A, Andreoli A, Puija A, Wang J, Pierson RN (1998): Total body potassium in healthy italians and americans. A cross-calibration study. In: *In vivo body composition studies*. Yasumura S, Wang J, Pierson RN (Eds), *Ann N Y Acad Sci*, **904**:366-368.

De Lorenzo A, Andreoli A, Battisti P, Talluri T, Yasumura S (1998): Total body capacitance correlates with total body potassium. In: *In vivo body composition studies*. Yasumura S, Wang J, Pierson RN (Eds), *Ann N Y Acad Sci*, **904**:259-262.

De Lorenzo A, Andreoli A, Deurenberg P (1997): Impedance ratio as a measure of water shifts. *Ann Nutr Met*, **41**:22-28.

De Lorenzo A, Andreoli A, Matthie J, Withers P (1997): Predicting body cell mass with bioimpedance by using theoretical methods: a technological review. *J Appl Physiol*, **82**(5):1542-1558.

De Lorenzo A, Candeloro N, Andreoli A, Deurenberg P (1995): Determination of intracellular water by multifrequency bioelectrical impedance. *Ann Nutr Met*, **39**:177-184

De Lorenzo A, Sasso GF, Andreoli A, Sorge R, Candeloro N, Cairella M (1995): Improved prediction formula for total body water assessment in obese women. *Int J Obes*, 535-538.

De Lorenzo A, Candeloro N, Bertini I, Talluri T, Pierangeli L (1995): Total body capacity correlated with basal metabolic rate. *Appl Radiat Isot*, **49**(5/6):493-494.

- De Lorenzo A, Deurenberg P, Andreoli A, Sasso GF, Palestini M, Docimo R (1994): Multifrequency impedance in the assessment of body water losses during dialysis. *Renal Physiology*, **17**:326-332.
- De Lorenzo A, Sorge SP, Iacopino L, Andreoli A, de Luca P, Sasso GF (1994): Fat free mass by bioelectrical impedance vs dual-energy X-ray Absorptiometry. *Appl Radiat Isot*, **49**(5/6):739-741.
- Demura Sh, Sato S and Kitabayashi T (2004): Percentage of total body fat as estimated by three automatic bioelectrical impedance analyzers. *J Physiol Anthropol Appl Human Sci*, **23** (3):93-99.
- Deurenberg P, Deurenberg-Yap M (2003): Validity of body composition methods across ethnic population groups. *Acta Diabetol*, **40**:S246–S249.
- Deurenberg P, Deurenberg-Yap M, Schouten FJ (2002): Validity of total and segmental impedance measurements for prediction of body composition across ethnic population groups. *Eur J Clin Nutr*, **56**:214–20.
- Deurenberg P, Andreoli A, de Lorenzo A (1996): Multi-frequency bioelectrical impedance: A comparison between the Cole-Cole modelling and Hanai equations with the classical impedance index approach. *Annals Hum Biol*, **23**(1):31-40.
- Deurenberg P, Schutz Y (1995): Body composition: Overview of methods and future directions of research. *Ann Nutr Metab*, **39**:325-333.
- Deurenberg P, Wolde-Gebriel Z, Schouten FJ (1995): Validity of predicted total body water and extracellular water using multifrequency bioelectrical impedance in an Ethiopian population. *Ann Nutr Metab*, **39**:234–41.
- Deurenberg P, van Malkenhorst E, Schoen T (1995): Distal vs proximal electrode placement in the prediction of total body water and extracellular water from multifrequency bioelectrical impedance. *Am J Hum Biol*, **7**:77-83.
- Deurenberg P, Broekhoff C, Andreoli A, de Lorenzo A (1994): The use of multi-frequency impedance in assessing changes in body water compartments. *Age & Nutr*, **5**(2):137-141.
- Deurenberg P, Weststrate JA, Seidell JC (1991): Body mass index as a measure of body fatness: age- and sex-specific prediction formula. *Brit J Nutr*, **65**:105–14.
- Deurenberg P, van der Kooy K, Leenen R, Weststrate J A, Seidell JC (1991): Sex and age specific prediction formulas for estimating body composition from bioelectrical impedance: a cross-validation study. *Int J Obes*, **15**:17-25.
- De Palo T, Messina G, Edefonti A, Perfumo F, Pisanello L, Peruzzi L, Piccoli A (2000): Normal values of the bioelectrical impedance vector in childhood and puberty. *Nutrition*, **16**(6):417-24.
- DiIorio BR, Scalfi L, Terracciano V, Bellizzi V (2004): A systematic evaluation of bioelectrical impedance measurement after hemodialysis session. *Kidney Int*, **65**:2435–2440.
- Di Iorio B, Bellizzi V (2000): Association of mortality and morbidity with bioimpedance analysis. *Kidney Int*, **58**:464-465.
- Downs JH, Haffjee A (1998): Nutritional assessment in the critically ill. *Curr Opin Clin Nutr Metab Care*, **1**:275-279.
- Dumler F (2003): Hypoalbuminemia is a marker of overhydration in chronic maintenance patients on dialysis. *ASAIO Journal*, 282-286.
- Dumler F, Kilates C (2000): Use of bioelectrical impedance techniques for monitoring nutritional status in patients on maintenance dialysis (Review). *Journal of Renal Nutrition*, **10**(3):116-124.

- Dumler F, Falla P, Butler R (1998): Impact of peritoneal dialysis modality and acidosis on nutritional status in peritoneal dialysis patients. *Adv. Peritoneal Dialysis*, **14**: 205-208.
- Dumler F (1997): Use of bioelectrical impedance analysis and dual-energy X-ray absorptiometry for monitoring the nutrition status of dialysis patients. *ASAIO Journal*, 256-260.
- Egger NG, Carlson GL, Shaffer JL (1999): Nutritional status and assessment of patients on home parenteral nutrition: anthropometry, bioelectrical impedance or clinical judgment. *Nutrition* **15**(1):1-6.
- Ellis KJ (2000) Human body composition: in vivo methods. *Physiol Rev*, **80**:649-680.
- Ellis KJ, Bell SJ, Chertow GM, Chumlea WC (1999): Bioelectrical impedance methods in clinical research: A follow-up to the NIH technology assessment conference. *Nutrition*, **15** (11, 12): 874-880.
- Ellis KJ, Shypailo RJ, Wong WW (1999): Measurement of body water by multifrequency bioelectrical impedance spectroscopy in a multiethnic pediatric population. *Am J Clin Nutr*, **70**:847-853.
- Ellis KJ, Wong WW (1998): Human hydrometry: comparison of multifrequency bioelectrical impedance with deuterium and bromine dilution. *J Appl Physiol*, **85**:1056.
- Evans WD, McClagish H, Trudgett C (1998): Factors affecting the *in vivo* precision of bioelectrical impedance analysis. *Appl Radiat Isot*, **49**(5/6):485-487.
- Fein P, Gundumalla G, Jordan A (2002): Usefulness of bioelectrical impedance analysis in monitoring nutrition status and survival of peritoneal dialysis patients. *Advances in Peritoneal Dialysis*, **18**:195-199.
- Fein A, Grossman RF, Jones JG, Goodman PC (1979): Evaluation of transthoracic electrical impedance in the diagnosis of pulmonary edema. *Circulation*, **60**(5):1156-1160.
- Fenech M, Jaffrin M (2004): Extracellular and intracellular volume variations during postural change measured by segmental and wrist-ankle bioimpedance spectroscopy. *IEEE Trans. Biom. Eng*, **51**(1): 166-175.
- Foster KR, Lukaski HC (1996): Whole body impedance – what does it measure? *Am J Clin Nutr*, **64**(suppl):338S-396S
- Foster KR, Schwan HP (1996): Dielectric properties of tissues. *Handbook of Biological Effects of Electromagnetic Fields*, Chapter 1; 25-103.
- Fogelholm M, van Marken Lichtenbelt (1997): Comparison of body composition methods: a literature analysis. *Eur J Clin Nutr*, **51**:495-503.
- Frerichs I, Hahn G, Schröder T, Hellige G (1998): Electrical impedance tomography in monitoring experimental lung injury. *Intensive Care Med*, **24**:829-836.
- Fuller NJ, Sawyer MB, Elia M (1994): Comparative evaluation of body composition methods and predictions, and calculation of density and hydration fraction of fat-free mass, in obese women. *Int J Obes*, **18**:503-512.
- Fusaro M, Munaretto G, Urso M, Piccoli A (2001): Severe obesity in hemodialysis: the utility of bioimpedance vector analysis. *Nephrol Dial Transplant*, **16**: 1273-1276.
- Gauyton AC and Hall JE(2001): *Textbook of medical physiology* (10th ed). Philadelphia, Saunders, p 274-329.

- Geddes LA, Baker LE (1967): The specific resistance of biological material – a compendium of data for the biomedical engineer and physiologist. *Med & Biol Engng*, **5**:271-293.
- Ghosh S, Meister D, Cowen S, Hannan WJ, Ferguson A (1997): Body composition at the bedside. *Eur Jour Gastroenterol Hepatol*, **9**:783-788.
- Goodman LR (1996): Congestive Heart Failure and Adult Respiratory Distress Syndrome: New Insights Using Computed Tomography. *Radiol Clin North Am*, **34**:33.
- Goran MI, Toth MJ, Poehlman ET (1997): Cross-validation of anthropometric and bioelectrical resistance prediction equations for body composition in older people using the 4-compartment model as a criterion method. *Am Geriatr Soc*, **45**:837-843.
- Grimnes S, Martinsen ØG (2005): Cole Electrical Impedance Model-A Critique and an Alternative. *IEEE Trans. Biomed. Eng*, **52**: 132-135.
- Grimnes S, Martinsen ØG and Schwan HP (2002): Interface phenomena and dielectric properties of biological tissue. *Encyclopedia of Surface and Colloid Science*. 2643-2652
- Grimnes S, Martinsen ØG (2000): *Bioimpedance and bioelectricity basics*. London, Academic Press.
- Gualdi-Russo E, Toselli S, Squintani (1997)L: Remarks on methods for estimating body composition parameters: reliability of skinfold and multiple frequency bioelectric impedance methods. *Z Morph Anthropol*, **81**(3):321-331.
- Guida B, Laccetti R, Trio R, Procino A (2005): Body mass index and survival differences in dialysis patients. (Letters to the Editor). *Kidney Int*, **67**:1634-1637.
- Guida B, De Nicola L, Trio R (2000): Comparison of vector and conventional bioelectrical impedance analysis in the optimal dry weight prescription in hemodialysis. *Am J Nephrol*, **20**:311-318.
- Guo SS, Wisemandle WA, Tyleshevsky FE (1997): Inter-machine and inter-method differences in body composition measures from dual energy X-ray absorptiometry. *Age & Nutr*, **8**(1):12-21.
- Guo SS, Chumlea WC, Cockram DB (1996): Use of statistical methods to estimate body composition. *Am J Clin Nutr*, **64**(3S):428S-435S.
- Haemrich D, Staelin ST, Tsai JZ (2003): In vivo electrical conductivity of hepatic tumours. *Physiol. Meas*, **24**:251-260.
- Hanai T (1968): Electrical properties of emulsions. In emulsion Science. London Academic Press.
- Hannan WJ, Cowen SJ, Plester C, Fearon KCH (1998): Proximal and distal measurements of extracellular and total body water by multi-frequency bio-impedance analysis in surgical patients. *Appl Radiat Isot*, **49**(5/6):621-622.
- Hannan WJ, Cowen SJ, Plester CE, Fearon KCH, deBeau A (1995): Comparison of impedance spectroscopy and multi-frequency bio-impedance analysis for the assessment of extracellular and total body water in surgical patients. *Clin Scien*, **89**:651-658.
- Hannan WJ, Cowen SJ, Fearon KCH, Plester CE, Falconer JS, Richardson RA (1994): Evaluation of multi-frequency bio-impedance analysis for the assessment of extracellular and total body water in surgical patients. *Clin Scien*, **86**:479-485.
- Heymsfield SB, Nuñez C, Testolin C, Gallagher D (2000): Anthropometry and methods of body composition measurement for research and field application in the elderly. *Europ J Clin Nutr*, **54**(Suppl 3):S26-S32.

- Heymsfield SB, Gallagher D, Grammes J, Nuñez C, Wang Z, Pietrobelli (1998): Upper extremity skeletal muscle mass: potential of measurement with single frequency bioimpedance analysis. *Appl Radiat Isot*, **49**(5/6):473-474.
- Heymsfield SB, Wang ZM, Baumgartner RN, Ross R (1997): Human body composition: advances in models and methods. *Annu Rev Nutr*, **17**:527-528.
- Heymsfield SB, Wang ZM, Visser M (1996): Techniques used in the measurement of body composition: an overview with emphasis on bioelectrical impedance analysis. *Am J Clin Nutr* **64**(Suppl):478S-484S.
- Heymsfield SB, Matthews D (1994): Body composition: research and clinical advances – 1993 ASPEN Research Workshop. *Jour Parent Enter Nutr*, **18**:91-103.
- Heyward VH (1998): Practical body composition assessment for children, adults and older adults. *Int Jour Sport Nutr*, **8**:285-307.
- Heyward VH (1996): Evaluation of body composition. Current issues. *Sports Med*, **22**(3):146-156.
- Hodgdon JA, Friedl KE, Beckett MB, Westphal KA, Shippee RL (1996): Use of bioelectrical impedance analysis measurements as predictors of physical performance. *Am J Clin Nutr*, **64**(suppl):463S-468S.
- Horstkotte D, Schulte HD, Niehues R (1993): Diagnostic and therapeutic considerations in acute, severe mitral regurgitation: Experience in 42 consecutive patients entering the intensive care unit with pulmonary edema. *J Heart Valve Dis*, **2**(5):512-522.
- Hotelling, H (1947): Multivariate Quality Control. In C. Eisenhart, M. W. Hastay, and W. A. Wallis, eds. *Techniques of Statistical Analysis*. New York: McGraw-Hill.
- Houtkooper LB, Lohman TG, Going SB, Howell WH (1996): Why bioelectrical impedance analysis should be used for estimating adiposity. *Am J Clin Nutr*, **64**:436S-448S.
- Hume R, Weyers E (1971): Relationship between total body water and surface area in normal and obese subjects. *J Clin Pathol*, **24**:234-238.
- Ikizler TA, Wingard RL, Harvell J (1999): Association of morbidity with markers of nutrition and inflammation in chronic hemodialysis patients: a prospective study. *Kidney Int*, **55**:1945-1951.
- Ishibe SH, Peixoto AJ (2004): Methods of assessment of volume status and intercompartmental fluid shifts in hemodialysis patients: Implications in clinical practice. *Seminars in Dialysis*, 37-43.
- Ishihara H, Suzuki A, Okawa H (2001): Comparison of initial distribution volume of glucose and plasma volume in thoracic fluid-accumulated patients. *Crit Care Med*, **29**(8):1532-1538.
- Jacobs DO (1996): Use of bioelectrical impedance analysis measurements in the clinical management of critical illness. *Am J Clin Nutr*, **64**(3S):498S-502S
- Jaffrin MY, Fenech M, de Fremont JF (2002): Continuous monitoring of plasma, interstitial, and intracellular fluid volumes in dialyzed patients by bioimpedance and hematocrit measurements. *ASAIO J*, 326-333.
- Jaffrin MY, Maasrani M, Gourrier A Le, Boudailliez B (1997): Extra-and intracellular volume monitoring by impedance during hemodialysis using Cole-Cole extrapolation. *Med Biol Eng Comput*, **35**:266-270.
- Jaffrin MY, Maasrani M, Boudailliez B (1996): Extracellular and intracellular fluid volume monitoring during dialysis by multiplequency impedancemetry. *ASAIO J*, **42**:M533-M538.

- Janssen I, Heymsfield SB, Baumgartner RN, Ross R (2000): Estimation of skeletal muscle mass by bioelectrical impedance analysis. *J Appl Physiol*, **89**:465-471.
- Jebb SA, Cole TJ, Doman D, Murgatroyd PR, Prentice AM (2000): Evaluation of the novel Tanita body fat analyzer to measure body composition by comparison with a four-compartment model. *Br J Nutr*, **83**:115-120.
- Johansson AN, Samuelsson O, Attman PO (2001): Limitations in anthropometric calculations of total body water in patients on peritoneal dialysis. *J Am Soc Nephrol*, **12**(3):568-573.
- Jolicoeur P (1999): Introduction to biometry. New York, Kluwer Academic/Plenum Publisher
- Jones CH, Smye SW, Newstead CG (1998): Extracellular fluid determined by bioelectric impedance and serum albumin in CAPD patients. *Nephrol Dial Transplant*, **13**: 393-397.
- Kanai H, Haeno M, Sakamoto K (1982): Electrical measurement of fluid distribution in legs and arms: Estimation of extra_cellular and intr_cellular fluid. *IEEE Frontiers of Engineering in Health Care*, 273-276.
- Kanetaka T (1990): Diagnosis of a special health check using Mahalanobi's Generalized Distance. *The ASI Journal*, **3**: 1.
- Khandelwal M, Kotharis J, Krishnan M (2003): Volume expansion and sodium balance in peritoneal dialysis patients. Part II: Newer insights in management. *Adv Perit Dial*, **19**:44-52.
- Kerkhof van J, Konings CJ, Krell van B (2003): Bioimpedance analysis and assessment of intracellular water in peritoneal dialysis patients. *Perit Dial Int*, **23**:591-605.
- Keller G, Blumberg A (1977): Monitoring of pulmonary fluid volume and stroke volume by impedance cardiography in patients on hemodialysis. *Chest*, **72**(1):56-62.
- Khan M, O'Hara R, Pohlman RL, Goldstein DB and Guha SK (2005): Multi-dimension applications of bioelectrical impedance analysis. *ASEP J*, **8**(1): 56-71.
- Klassen P, Mazariegos M, Deurenberg P, Solomons NW, Furst P (2000): Hydrational status assessed by bioelectrical impedance spectroscopy and dilution methods in patients with classical Dengue fever. *Annals NY Acad Sci*, **904**:163-170
- Ketteler M, Gross ML and Ritz E (2005): Calcification and cardiovascular problems in renal failure. *Kidney Int*, **67**:S120-S127.
- Koc M, Richards HB, Bihorac A, Ross EA, Schold JD and Segal M (2005): Circulating endothelial cells are associated with future vascular events in hemodialysis patients. *Kidney Int*, **67**:1078-1083.
- Kojima M, Hasegawa Y (1994): Prediction of urinary continence recovery among patients with brain diseases using Mahalanobis Distance, standarization and quality control. *The ASI Journal*, **3**: 47.
- Kooman JP, Leunissen KML, Luik AJ (1998): Salt and hypertension in end-stage renal disease. *Blood Purif*, **16**:301-311.
- Kööbi T, Kähönen M, Koskinen M, Kaukinen S, Turjanmaa VMH (2000): Comparison of bioimpedance and radioisotope methods in the estimation of extracellular water volume before and after coronary artery bypass grafting operation. *Clin Physiol*, **20**(4):283-291.

- Konings CJ, Kooman IP, Schonck M (2003): Influence of fluid status on techniques used to assess body composition in peritoneal dialysis patients. *Perit Dial Int*, **23**:184-190.
- Konings CJ, Kooman IP, Schonck M (2002): Assessment of fluid status in peritoneal dialysis patients. *Perit Dial Int*, **22**:683-692
- Koizumi N, Warisawa S, Hashizume H (2003) : Impedance controller and its clinical use of the remote ultrasound diagnostic system. Proceedings of the *IEEE International Conference on Robotic & Automation*, Taipei, 676-683.
- Kotler DP, Burastero S, Wang J, Pierson RN (1996): Prediction of body cell mass, fat-free mass and total body water with bioelectrical impedance analysis: effects of race, sex and disease. *Am J Clin Nutr*, **64**(3 suppl):489S-497S
- Kobert MS (1998): Evaluation of ultrafiltration failure. *Adv Ren Replace Ther*, **5**:194-204.
- Kuczmarski RJ (1996): Bioelectrical impedance analysis measurements as part of a national nutrition survey. *Am J Clin Nutr*, **64**(suppl):453S-458S
- Kuhlmann MK, Zhu F, Seibert E, Levin NW (2005): Bioimpedance, dry weight and blood pressure control: new methods and consequences. *Curr Opin Nephrol Hypertens*, **14**:543-549
- Kushner RF, Gudivaka R, Schoeller DA (1996): Clinical characteristics influencing bioelectrical impedance analysis measurements. *Am J Clin Nutr*, **64**(3S):423S-427S.
- Kushner RF, de Vries PMJM, Gudivaka R (1996): Use of bioelectrical impedance analysis measurements in the clinical management of patients undergoing dialysis. *Am J Clin Nutr*, **64**(Suppl):503S-509S.
- Kushner RF, Schoeller DA, Fjeld CR (1992): Is the impedance index (ht^2/R) significant in predicting total body water? *Am J Clin Nutr*, **56**:835-839.
- Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, Heitmann BL, Kent-Smith L, Melchior JC, Pirlich M, Scharfetter H, Schols AMWJ, Pichard C (2004): Bioelectrical impedance analysis- Part II: Utilization in clinical practice. *Clin Nutr*, **23**:1430-1453.
- Kyle UG, Pichard C (2000): Dynamic assessment of fat-free mass during catabolism and recovery. *Curr Opin Clin Nutr Metab Care*, **3**:317-322.
- Kyle UG, Pichard C, Rochat T, Slosman DO, Fitting JW, Thiebaud D (1998): New bioelectrical impedance formula for patients with respiratory insufficiency: comparison to dual-energy X-ray absorptiometry. *Eur Respir J*, **12**:960-966.
- Lameire N, Van Biesen W (2004): Hypervolemia in peritoneal dialysis patients Review. *J Nephrol*. **17**(8S): S58-66.
- Lederman SA, Paxton A, Heymsfield SB, Wang J, Thornton J, Pierson RN (1997): Body fat and water changes during pregnancy in women with different body weight and weight gain. *Obst Gynecol*, **90**(4):483-488.
- Lee SW, Song JH, Kim GA (2001): Assessment of total body water from anthropometry-based equations using bioelectrical impedance as reference in Korean adult control and hemodialysis subjects. *Nephrol Dial Trasplantation*, **16**:91-97.
- Leiter LA, Lukaski HC, Kenny DJ, Barnie A, Camelon K, Sie R (199): The use of bioelectrical impedance analysis (BIA) to estimate body composition in the diabetes control and complications trial (DCCT). *Int J Obesity* **18**:829-835.

Lentner C (1982): Introduction to statistics. Statistical tables. Mathematical formulae, in Geigy Scientific Tables (vol 2, 8th ed), Basle, Ciba-Geigy Limited.

Liang MTC, Su HF, Lee NY (2000): Skin temperature and skin blood flow affect bioelectric impedance study of female fat free mass. *Med Sci Sports Exer* 32(1):221-227.

Lopot F, Nejedly B, Novotna H (2002): Age-related extracellular to total body water volume ratio (ECV/TBW) –Can it be used for “dry weight” determination in dialysis patients? Application of multifrequency bioimpedance measurement. *J of Artif. Kidney and Dial*, 25(8):762-769.

Lozano A, Rosell J and Pallàs-Areny R (1995): A Multifrequency multichannel electrical impedance data acquisition system for body fluid shift monitoring. *Physiol Meas*, 16:227-237.

Lozano A, Rosell J, Pallàs-Areny R (1995): Errors electrical impedance measurements due to electrode repositioning and postural changes. *Physiol Meas*, 16:121-130.

Lozano A, Rosell J, Riu PJ and Pallàs-Areny R (1995): Segmental body fluid shift estimation during HDT positions by electrical impedance measurements. Proceedings of the IX International Conference on Electrical Bio-Impedance. Heidelberg (Germany), 233-236.

Lozano A, Rosell J, Riu PJ and Pallàs-Areny R (1994): Multifrequency multisegment electrical impedance measurements for hemodialysis control. Proceedings of the V International Symposium on Biomedical Engineering. Santiago de Compostela (España), 213-214.

Lozano A, Riu PJ, Rosell J and Pallàs-Areny R (1992): Multifrequency segmental electrical impedance measurements for body fluid shift determination. Proceedings of the 8th. International Conference on Electrical Bio-impedance. Kuopio (Finland), 115-117.

Lukaski HC, Siders WA (2003): Validity and accuracy of regional bioelectrical impedance devices to determine whole-body fatness. *Nutrition*, 19: 851-857.

Lukaski HC (1999): Requirements for clinical use of bioelectrical impedance analysis (BIA). Electrical Bioimpedance Methods: Applications to Medicine And Biotechnology. *Annals of the New York Academy of Sciences*, 873:72-76.

Lukaski HC (1996): Biological indexes considered in the derivation of the bioelectrical impedance analysis. *Am J Clin Nutr*, 64:397S-404S.

Lukaski HC, Scheltinga RM (1994): Improved sensitivity of the tetrapolar bioelectrical impedance method to assess fluid status and body composition: use of proximal electrode placement. *Age Nutr*, 5:123-129.

Lukaski HC, Bolonchuck WW, Hall CB, Siders WA (1986): Validation of tetrapolar bioelectrical impedance method to assess human body composition. *J Appl Physiol*, 60:1327-1332.

Maasrani M, Jaffrin MY, Boudailliez (1997): Continuous measurements by impedance of haematocrit and plasma volume variations during dialysis. *Med Biol Eng Comput*, 167-171.

Madore F, Wuest M, Ethier JH (1994): Nutritional evaluation of hemodialysis patients using an impedance index. *Clin Nephrol*, 41(6):377-382.

Maggiore Q, Nigrelli S, Ciccarelli C (1996): Nutritional and prognostic correlates of bioimpedance indexes in hemodialysis patients. *Kidney Int*, 50:2103-2108.

Mahalanobis, PC (1930): On Tests and Measures of Groups Divergence I: *Journal of the Asiatic Society of Benegal*, 26: 541.

- Mallamaci F, Tripepi G, Cutrupi S (2005): Prognostic value of combined used of biomarkers of inflammation, endothelial dysfunction, and myocardial pathology in patients with ESRD. *Kidney Int*, **67**:2330-2337.
- Matthay, MA (1995): Acute hypoxemic respiratory failure: Pulmonary edema and ARDS. In: George, RB, Light, RW, Matthay, MA, et al (Eds), 3rd ed, Chest Medicine. Essentials of Pulmonary and Critical Care Medicine, Williams & Wilkins, Baltimore, 593.
- Mazariegos M, Klassen P, Solomons NW, Furst P (2000): Bioelectrical impedance spectroscopy in health and disease: Correspondence between whole body and segmental bioelectrical impedance spectroscopy indices in patients with classical Dengue fever. *Annals NY Acad Sci*, **904**:205-209.
- Mayer M, Brunner P, Merwa R and Scharfetter H (2005): Monitoring of lung edema using focused impedance spectroscopy: a feasibility study. *Physiol Meas* **26**:185-192.
- McCullough PA (2005): Evaluation and treatment of coronary artery disease in patients with end-stage renal disease. *Kidney Int*, **67**:S51-S58.
- Morrison DF (1967): Multivariate Statistical Methods. New York, MacGraw Hill.
- Montgomery LD, Parmet AJ, Booher CR (1993): Body volume changes during simulated microgravity: Auditory changes, segmental fluid redistribution, and regional hemodynamics. *Annals of Biomedical Engineering*, **21**:417-433.
- Mushnick R, Fein PA, Mittman N (2003): Relationship of bioelectrical impedance parameters to nutrition and survival in peritoneal dialysis patients. *Kidney Int*, **64**:S53-S56.
- Nanovic L (2005): Electrolytes and fluid management in hemodialysis and peritoneal dialysis. *Nutr Clin Pract*, **20**: 192-201.
- National kidney foundation (2001): K/DOQI clinical practice guidelines for peritoneal dialysis adequacy. *Am J Kidney Dis*, **37** (Suppl): 65S-136S.
- Newell JC, Ren X, Larson-Wiseman JL, Danyleiko MD (1996): Assessment of acute pulmonary lung edema in dogs by electrical impedance imaging. *IEEE Trans Biom Eng*, **43**(2):133-138.
- Nierman DM, Eisen DI, Fein ED (1996): Transthoracic bioimpedance can measure extravascular lung water in acute lung injury. *J Surgical Research*, **65**:101-108.
- Noble TJ, Harris ND, Morice AH (2000): Diuretic induced changes in lung water assessed by electrical impedance tomography. *Physiol Meas* **21**:155-163.
- Noble TJ, Morice AH, Milnes P (1996): Electrical impedance tomographic spectroscopy in the detection of increased lung water in heart failure. *18th Annual International Conference IEEE Engineering in Medicine and Biology Society*, 3.4.2: Applications of Electrical Impedance Tomography.
- Noordegraaf AV, van der Meer BJM, de Vries JPPM, de Vries PMJM (1995): Determination of the relation between alterations of total body water and thoracic fluid content during ultrafiltration by bioelectrical impedance analysis. *Nephrol Dial Transplant*, **10**:382-385.
- Núñez C, Gallagher D, Visser M, Pi-Sunyer FX, Wang Z, Heymsfield SB (1997): Bioimpedance analysis: evaluation of leg-to-leg system based on pressure contact foot-pad electrodes. *Med Sci Sports Exerc*, **29**(4):524-531.
- Oldham NM (1996): Overview of bioelectrical impedance analyzers. *Am J Clin Nutr*, **64**(3S):405S-412S.
- Ojanen S, Koivisto AM, Kööbi T (2002): Isolated ultrafiltration affects dynamic vectorcardiographic ischemia monitoring parameters: *Clinical Nephrology*, **57**:359-364.

- Ojanen S, Kööbi T, Koivisto AM (2002): Hemodialysis causes changes in dynamic vectorcardiographic ischemia monitoring parameters: *Clinical Nephrology* 2002; 54:227-233
- Ott M, Fischer H, Polat H (1995): Bioelectrical impedance analysis as a predictor of survival in patients with human immunodeficiency virus infection. *J Acquired Immune Deficiency Syndromes Human Retrovirology*, 9:20-25.
- Ott M, Lembcke B, Fischer H (1993): Early changes of body composition in human immunodeficiency virus-infected patients: tetrapolar body impedance analysis indicates significant malnutrition. *Am J Clin Nutr*, 57:15-19.
- Organ LW, Bradham B, Gore DT (1994): Segmental bioelectric impedance analysis: theory and application of a new technique. *J Appl Physiol*, 77:98-112.
- O'Riordan E, Chen J, Brodsky SV, Smirnova I, Hong Li (2005): Endothelial cell dysfunction: The syndrome in making. *Kidney Int*, 67:1654-1658
- Paniagua R, Amato D, Vonesh E (2002): Effects of increased peritoneal clearances on mortality rates in peritoneal dialysis: ADEMEX, a prospective, randomized, controlled trial. *J Am Soc Nephrol*, 13:1307-1320.
- Passadakis P, Sud K, Dutta A (1999): Bioelectrical impedance analysis in the evaluation of the nutritional status of continuous ambulatory peritoneal dialysis patients. *Adv Perit Dial*, 15:147-152.
- Patel RV, Peterson EL, Silverman N, Zarowitz BJ (1996): Estimation of total body and extracellular water in post-coronary artery bypass graft surgical patients using single and multiple frequency bioimpedance. *Crit Care Med*, 24(11):1824-1828.
- Patel RV, Matthie JR, Withers PO, Peterson EL, Zarowitz BJ: Estimation of total body water and extracellular water using single and multiple-frequency bioimpedance. *Ann Pharmacother* 1994; 28:565-569
- Peña D (2001): Fundamentos de estadística. Ciencias Sociales. Alianza Editorial.
- Piccoli A (2005): Whole Body-Single frequency bioimpedance. *Contrib Nephrol Basel Karger*, 49: 150-161.
- Piccoli A, Pastori G, Guizzo M (2005): Equivalence of information from single versus multiple frequency bioimpedance vector analysis in hemodialysis. *Kidney Int*, 67:301-313.
- Piccoli A for the Italian HD-BIA Study Group (2004): Bioimpedance vector migration up to three days after the hemodialysis session. *Kidney Int*, 66:2088-2094.
- Piccoli A, for the Italian CAPD-BIA study group (2004): Bioelectrical impedance vector distribution in peritoneal dialysis patients with different hydration state. *Kidney Int*, 65:1050-1063.
- Piccoli A (2002): Patterns of bioelectrical impedance vector analysis: Learning from electrocardiography and forgetting electric circuit models. *Nutrition*, 18: 520-521
- Piccoli A, Pillon L, Dumler F (2002): Impedance vector distribution by sex, race, body mass index, and age in the United States: standard reference intervals as bivariate Z scores. *Nutrition*, 18:153-167.
- Piccoli A, Nescolarde L, Rosell J (2002): Análisis Convencional y Vectorial de Bioimpedancia en la Práctica Clínica. *Revista Española de Nefrología*, XXII: 230-240.
- Piccoli A, Pastori G (2002): *BIVA software*. University of Padova.

- Piccoli A, Pittoni G, Facco E (2000): Relationship between central venous pressure and bioimpedance vector analysis in critically ill patients. *Crit Care Med*, **28**:132-137.
- Piccoli A, Pillon L, Tabbi MG (1999): Major confounders for reactance as a marker of malnutrition in hemodialysis patients. *Kidney Int*, **56**:2311-2312.
- Piccoli A, Pillon L, Pisanello L (1999): Electrical maturation trajectory of human tissues identified by bioelectrical impedance vector analysis. *Nutrition*, **15**:77-78.
- Piccoli A, Brunani A, Savia G (1998): Discriminating between body fat and fluid changes in the obese adult using bioimpedance vector analysis. *Int J Obesity*, **22**:97-104.
- Piccoli A, for the Italian HD-BIA Study Group (1998): Identification of operational clues to dry weight prescription in hemodialysis using bioimpedance vector analysis. *Kidney Int*, **53**:1036-1043.
- Piccoli A, Pillon L, Favaro E (1997): Asymmetry of the total body water prediction bias using the impedance index. *Nutrition*, **13**:438-441.
- Piccoli A, Piazza P, Noventa D (1996): A new method for monitoring hydration at high altitude by bioimpedance analysis. *Med Sci Sports Exerc*, **28**:1517-1522.
- Piccoli A, Rossi B, Pillon L (1996): Body fluid overload and bioelectrical impedance analysis in renal patients. *Miner Electrolyte Metab*, **22**:76-78.
- Piccoli A, Nigrelli S, Caberlotto A, Bottazzo S, Rossi B, Pillon L, Maggiore Q (1995): Bivariate normal values of the bioelectrical impedance vector in adult and elderly populations. *Am J Clin Nutr*, **61**:269-270.
- Piccoli A, Rossi B, Pillon L, Bucciantè G (1994): A new method for monitoring body fluid variation by bioimpedance analysis: The RXc graph. *Kidney Int*, **46**:534-539.
- Piccoli A, Rossi B, Pillon L (1994): Operational equivalence between segmental and whole-body bioelectrical impedance in renal patients. *Am J Clin Nutr*, **59**:675-676
- Pichard C, Kyle UG, Bracco D, Slosman DO, Morabia A, Schutz Y (2000): Reference values of fat-free and fat masses by bioelectrical impedance analysis in 3393 healthy subjects. *Nutrition*, **16**:245-254.
- Pichard C, Kyle UG, Slosman DO (1999): Fat-free mass in chronic illness: Comparison of bioelectrical impedance and dual-energy X-Ray absorptiometry in 480 chronically ill and healthy subjects. *Nutrition*, **15**(9):668-676.
- Pichard C, Kyle UG (1998): Body composition measurements during wasting diseases. *Curr Opin Clin Nutr Metab Care*, **1**:357-361.
- Pichard C, Kyle UG, Gremion G, Gerbase M, Slosman DO (1997): Body composition by x-ray absorptiometry and bioelectrical impedance in female runners. *Med Sci Sports Exerc*, **29**(11):1527-1534.
- Pierson RN, Wang J, Thornton JC (1997): Measurement of body composition: Applications in hormone research. *Horm Res*, **48**(suppl 1):56-62.
- Pietrobelli A, Rubiano F, St-Onge MP, Heymsfield SB (2004): New bioimpedance analysis: improved phenotyping with whole-body analysis. *Eur J Clin Nutr*, **58**:1479-1484.
- Pietrobelli A, Morini P, Battistini N, Chiumello, Nuñez C, Heymsfield SB (1998): Appendicular skeletal muscle mass: prediction from multiple frequency segmental bioimpedance analysis. *Eur J Clin Nutr*, **52**:507-511.

- Pietrobelli A, Wang Z, Heymsfield SB (1998): Techniques used in measuring human body composition. *Curr Opin Clin Nutr Metab Care*, **1**:439-448.
- Pillon L, Piccoli A, Lowrie EG (2004): Vector length as a proxy for the adequacy of ultrafiltration in hemodialysis. *Kidney Int*, **66**:1266-1271.
- Plum J, Schonike G, Kleophas W (2001): Comparison of body fluid distribution between chronic hemodialysis and peritoneal dialysis patients as assessed by biophysical and biochemical methods. *Nephrol Dial Transplant*, **16**:2378-2385.
- Raaijmakers E, Faes th JC, Kunst PWA (1998): The influence of extravascular lung water on cardiac output measurements using thoracic impedance cardiography. *Physiol Meas*, **19**:491-499.
- Raaijmakers E, Faes th JC, Meijer JM (1998): Estimation of non-cardiogenic pulmonary oedema using dual-frequency electrical impedance. *Med Biol Eng Comput*, **36**:461-466.
- Ramos MU, Labree JW, Remole W, Kubiceck WG (1995): Transthoracic electric impedance. *A clinical guide of pulmonary fluid accumulation in congestive heart failure*, 671- 676.
- Rallison LR, Kushner RF, Penn D, Schoeller DA (1993): Errors in estimating peritoneal fluid by bioelectrical impedance analysis and total body electrical conductivity. *J Am Coll Nutr*, **12**: 66-72.
- Ravaglia G, Forti P, Maioli F, Boschi F, Cicognani A, Gasbarrini G (1999): Measurement of Body Fat in healthy elderly men: A comparison of methods. *J Geront*, **54A**(2)M70-M76
- Rees AE, Ward LC, Cornish BH, Thomas BJ (1999): Sensitivity of multiple frequency bioelectrical impedance analysis to changes in ion status. *Physiol Meas*, **20**:349-362.
- Ritz E (2005): Renal dysfunction as a novel risk factor: Microalbuminuria and cardiovascular risk. *Kidney Int*, **67**:S25-S28.
- Riu P, Nescolarde L, Vidal A (2004): Effect of dielectric anisotropy of tissues in body composition analysis. Proceedings of the *XII International Conference on Electrical Bioimpedance*. Gdansk (Poland). ISBN-83-917681-6-3 pp 229-232.
- Rosell J, Casas O, Bragós R, Riu PJ and Andersen LJ (1997): Estimation of intra and extra-cellular segmental volumes using electrical impedance measurements Proceedings of the *World Congress on Medical Physics and Biomedical Engineering*. Nice (France). ISSN:0140-018, pp 33
- Rosell Ferrer J, Riu Costa P, Pallas Areny R, Elvira Canas J, Bragos Bardia R, NTE, S.A (2000): Apparatus and procedure for measuring volumes and global and segmental corporal composition in human beings. *US Patent office*. Patent number, US6151523.
- Rosell Ferrer J, Elvira Canas J, Riu Costa P, Bragos Bardia R, Pallas Areny R, NTE S.A (1998): Apparatus for measuring volumes and global and segmental corporal composition in human beings. *European Patent office*. Patent number, EP865763.
- Ross AN, Westendorp RGJ, Brand R, Souverijn JHM, Frölich M, Meinders AE (1995): Predictive value of tetrapolar body impedance measurements for hydration status in critically ill patients. *Intensive Care Med*, **21**:125-131.
- Roubenoff R, Baumgartner RN, Harris T (1997): Application of bioelectrical impedance analysis to elderly populations. *J Geront*, **52A**(3):M129-M136.
- Roubenoff R (1996): Application of bioelectrical impedance analysis for body composition to epidemiological studies. *Am J Clin Nutr*, **64**(suppl):459S-462S.
- Roubenoff R, Dallal GE, Wilson PWF (1995): Predicting body fatness: the body mass index vs estimation by bioelectrical impedance. *Am J Pub Health* **85**(5):726-728.

- Sadlier RJ, Fox RA (2001): Detection and quantification of intraperitoneal fluid using electrical impedance tomography. *IEEE Trans Biom Eng*, **48**:484-491.
- Sakamoto K, Kanai H, Furuya N (2004): Electrical admittance method for estimating fluid removal during artificial dialysis. *Med Biol Eng Comp*, **42**:356-365.
- Scalfi L, Marra M, Caldara A (1999): Changes in bioimpedance analysis after stable refeeding of undernourished anorexic patients. *Int J Obes*, **23**:133-137.
- Scharfetter H, Schlager T, Stollberger R, Felsberger R (2001): Assessing abdominal fatness with local bioimpedance analysis: Basics and experimental findings. *Int Journal of Obesity*, **25**(4):502-511.
- Scharfetter H, Hartinger P, Hinghofer-Szalkay H, Hutten H (1998): A model of artefacts produced by stray capacitance during whole body or segmental bioimpedance spectroscopy. *Physiol Meas*, **19**(2):247-61.
- Scharfetter H, Monif M, László Z, Lambauer T, Hutten H, Hinghofer-Szalkay H (1997): Effect of postural changes on the reliability of volume estimations from bioimpedance spectroscopy data. *Kidney Int*, **51**:1078-1087.
- Scharfetter H, Wirnsberger G, Hutten H (1994): Comparison of measured and simulated volume shifts between the intra and the extracellular space in dialysis patients. *Annual International Conference of the IEEE Engineering in Medicine and Biology*. **2**:1142-1143.
- Schmit R, Dumler F, Cruz C (1993): Indirect measures of total body water may confound precise assessment of peritoneal dialysis adequacy. *Perit Dial Int*, **13**:S224-S226.
- Schuster HP, Schuster CJ, Gilfrich HJ, Schölmerich (1980): Transthoracic electrical impedance during extracorporeal hemodialysis in acute respiratory failure ("Shocked Lung Syndrome"). *Intens Care Med*, **6**:147-154.
- Smak Gregoor PJH (2005): The differential impact of risk factors on mortality in hemodialysis and peritoneal dialysis. (Letters to the Editor). *Kidney Int*, **67**:2504-2507.
- Schwan HP (1957): Electrical properties of tissues and cell suspensions. In: Lawrence JH, Tobias CA (eds) *Advances in biological and medical physics*. Vol V, 147-209. Academic Press, New York.
- Simons JPFHA, Schols AMWJ, Westertep KR, ten Velde GPM, Wouters EFM (1995): The use of bioelectrical impedance analysis to predict total body water in patients with cancer cachexia. *Am J Clin Nutr*, **61**:741-745.
- Snedecor, GW, Cochran WG (1989): *Statistical Methods*, Eighth Edition, Iowa State University Press.
- Song JH, Lee SW, Kim GA and Kim MJ (1999): Measurement of fluid shift in CAPD patients using segmental bioelectrical impedance analysis. *Perit Dialysis Int*, **19**:386-390.
- Spiegel DM, Bashir K and Fisch B (2000): Bioimpedance resistance ratios for the evaluation of dry weight in hemodialysis. *Clinical Nephrology*, **53**(2):108-114.
- Stall SH, Ginsberg NS, DeVita MV (1996): Comparison of five body-composition methods in peritoneal dialysis patients. *Am J Clin Nutr*, **64**:125-130.
- Staub N, Hogg JC (1980): Conference report of a workshop on the measurement of lung water. *Crit Care Medicine*, **8**(12):752-759.
- Stolarczyk LM, Heyward VH, Van Loan MD, Hicks VL, Wilson WL, Reano LM (1997): The fatness-specific bioelectrical impedance analysis equations of Segal et al: are they generalizable and practical? *Am J Clin Nutr*, **66**:8-17.

- Stolarczyk LM, Heyward VH, Hicks VL, Baumgartner RN (1994): Predictive accuracy of bioelectrical impedance in estimating body composition of Native American women. *Am J Clin Nutr*, **59**:964-970.
- Stout JR, Eckerson JM, Housh TJ, Jonson GO, Betts NM (1994): Validity of percent body fat estimations in males. *Med Sci Sports Exerc*, **26**(5):632-636.
- Sun SS, Chumlea WC, Heymsfield SB (2003): Development of bioelectrical impedance analysis prediction equations for body composition with the use of a multicomponent model for use in epidemiologic surveys. *Am J Clin Nutr* **77**:331-40.
- Stucliffe JF (1996): A review of *in vivo* experimental methods to determine the composition of the human body. *Phys Med Biol*, **41**:791-833.
- Swan PD, McKonnell KE (1999): Anthropometry and bioelectrical impedance inconsistently predict fatness in women with regional adiposity. *Med Sci Sports Exerc*, **31**(7):1068-1075.
- Taguchi G (1987): System of Experimental Design, American Supplier Institute, Unipub.
- Tai ES, Ho SC, Fok ACK, Tan CE (1999): Measurement of obesity by anthropometry and bioelectrical impedance analysis: Correlation with fasting lipids and Insulin resistance in an Asian population. *Ann Acad Med Singapore*, **28**:445-450.
- Tan YX, Nuñez C, Sun Y, Zhang K, Wang Z, Heymsfield SB (1997): New electrode system for rapid whole-body and segmental bioimpedance assessment. *Med Sci Sports Exerc*, **29**(9):1269-1273.
- Tempel G, Jelen S, Hundelshausen BV (1988): Transthoracic electrical impedance in anaesthesia and intensive care. *Resuscitation*, **6**:97-105.
- Than N, Woodrow G, Oldroyd (2000): Effect of peritoneal fluid on whole body and segmental multiple frequency bioelectrical impedance in patients on peritoneal dialysis. *Eur J Clin Nutr*, **54**:450-451.
- Thomas BJ, Cornish BH, Ward LC, Jacobs A (1999): Bioimpedance: Is it a predictor of true water volume? *Annals of the New York Academy of Sciences*, **873**:89-93.
- Thomas BJ, Cornish BH, Ward LC, Patterson MA (1998): A comparison of segmental and wrist-to-ankle methodologies of bioimpedance analysis. *Appl Radiat Isot*, **49**(5/6):477-478.
- Thomas BJ, Ward LC, Cornish BH (1997): Bioimpedance spectrometry in the determination of body water compartments: Accuracy and clinical significance. *Appl Radiat Isot*, **49**: 447-455.
- Thompson CM, Kong CH, Lewis CA (1993): Can bio_electrical impedance be used to measured total body water in dialysis patients ? *Physiol Meas*, **14**:455-461.
- Tomashefski, JF (1990). Pulmonary pathology of the adult respiratory distress syndrome. *Clin Chest Med*, **11**:593.
- Toso S, Piccoli A, Gusella M, Menon D, Bononi A, Crepaldi G (2000): Altered tissue electric properties in lung cancer patients as detected by bioelectric impedance vector analysis. *Nutrition*, **16**(2):120-4.
- Turner AA, Lozano-Nieto A and Bouffard Marcel (2002): Comparison of segmental and global bioimpedance spectroscopy errors using generalizability theory. *Physiol Meas*, **23**:43-57.
- Tzamaloukas AH, Murata GH, Vanderjagt DJ, Glew RH (2001): Estimates of body water, fat-free mass, and body fat in patients on peritoneal dialysis by anthropometric formulas. *Kidney International*, **63**:1605-1617.

- Valensise H, Andreoli A, Lello S, Magnani F, Romanini C, de Lorenzo A (2000): Multifrequency bioelectrical impedance analysis in women with a normal and hypertensive pregnancy. *Am J Clin Nutr*, **72**:780-783.
- Van de Kerkhof J, Konings CJ, Van Kreel B (2003): Bioimpedance analysis and assessment of intracellular water in peritoneal dialysis patients. *Perit Dial Int*, **23**: 591-605.
- Van Loan MD, Strawford A, Jacob M and Hellerstein M (1999): Monitoring changes in fat-free mass in HIV-positive men with hypotestosteronemia and AIDS wasting syndrome treated with gonadal hormone replacement therapy. *AIDS*, **13**: 241-248.
- Van Loan MD (1998): Estimates of fat-free mass by densitometry, DEXA and bioimpedance spectroscopy in Caucasian and Chinese American women. *Appl Radiat Isot*, **49**(5/6):751-752.
- Van Loan MD (1997): Multicompartment model of body composition assessment in Chinese American adults. *Am J Hum Biol*, **9**:21-27.
- Van Loan MD, Kopp LE, King JC, Wong WW, Mayclin PL (1995): Fluid changes during pregnancy: use of bioimpedance spectroscopy. *J Appl Physiol*, **78**(3):1037-1042.
- Van Marken Lichtenbelt WD, Westerterp KR, Wouters L (1994): Validation of bioelectrical-impedance measurements as a method to estimate body-water compartments. *Am J Clin Nutr*, **60**:159-166.
- Visauta Vinacua B (2001): Análisis estadístico con SPSS para Windows. Tomo II. McGraw Hill.
- Wabel P, Moissl U, Thul P (2002): Defference between monofrequency bioimpedance analysis (BIA) and multifrequency bioimpedance spectroscopy (BIS)- a comparative study.
- Wagner DR, Heyward VH (2000): Measures of body composition in blacks and whites: A comparative review. *Am J Clin Nutr*, **71**(6):1392-1402.
- Wagner DR, Heyward VH (1999): Techniques of body composition assessment: a review of laboratory and field methods. *Research Quart Exerc Sport*, **70**(2):135-149.
- Wagner DR, Heyward VH, Kocina PS, Stolarczyk LM, Wilson WL (1997): Predictive accuracy of BIA equations for estimating fat-free mass of black men. *Med Sci Sports Exerc*, **29**:969.
- Ward LD, Heitman BL, Craig P (2000): Association between Ethnicity, Body Mass Index, and Bioelectrical Impedance. *Ann N Y Acad Sci*, **258**:199-202.
- Ward LC, Byrne NM, Rutter K, Hennoste L (1997): Reliability of multiple frequency bioelectrical impedance analysis : An Intermachine comparison. *Am. J of Human Biology*, **9**:63-72.
- Watson PE, Watson ID, Batt RD (1980): Total body water volumes for adult males and females estimated from simple anthropometric measurement. *Am J Clin Nutr*, **33**:27-29.
- Wilmore KM, McBride PJ, Wilmore JH (1994): Comparison of bioelectric impedance and near-infrared interactance for body composition assessment in a population of self-perceived overweight adults. *Int J Obes*, **18**:375-381.
- Winer BJ (1962): Statistical Principles in Experimental Design. New York: McGraw-Hill.
- Woo EJ, Hua P, Webster JG, Tompkins WJ (1992): Measuring lung resistivity using electrical I pedance tomography. . *IEEE Trans Biom Eng*, **39**(7):756-760.
- Woodrow G, Oldroyd B, Wright A (2004): Abnormalities of body composition in peritoneal dialysis patients. *Perit Dial Int*, **24**:169-175.

Woodrow G, Oldroyd B, Wright A (2003): Comparison of anthropometric equations for estimation of total body water in peritoneal dialysis patients. *Nephrol Dial Transplant*, **18**:384-389.

Woodrow G, Oldroyd B, Stables G (2000): Effects of icodextrin in automated peritoneal dialysis on blood pressure and bioelectrical impedance analysis. *Nephrol Dial Transplant*, **15**:862-866.

Wotton MJ, Thomas BJ, Cornish BH, Ward LC: Comparison of whole body and segmental bioimpedance methodologies for estimating total body water: *Annals NY Acad Sci* 2000: ISBN 1-57331-258-4 : 181-186

Wong KC, Xiong DW, Kerr PG (1995): Kt/V in CAPD by different estimations of V. *Kidney Int*, **48**:563-569.

Wong V, How TV, ward R (1994): Input impedance of radiocephalic arteriovenous fistulae for hemodialysis access: its value in predicting early failure: *Physiol Meas*, **15**:469-480.

Zaluska WT, Schneditz D, Swatowski A (2003): Comparison of prescribed and delivered doses of dialysis using anthropometrically and bioelectrically measured patients volumes. *Med Sci Monit* **9**(9):CR405-410.

Zaluska WT, Malecka T, Swatowski A (2002): Changes of extracellular volumes measured by whole and segmental bioimpedance analysis during hemodialysis in end-stage renal disease (ESRD) patients. *Annales Universitatis Marie Curie-Lublin, Polonia*. **2**(131): 337-341.

Zaluska WT, Schneditz D, Kaufman AM, Morris AT, Levin NW (1998): Relative underestimation of fluid removal during hemodialysis-hypotension measured by wrist to ankle bioimpedance. *ASAIO J*, **44**:823-827.

Zdolsek HJ, Lindahl OA, Sjöberg F (2000): Non-invasive assessment of fluid volume status in the interstitium after hemodialysis: *Physiol Meas*, **21**:211-220.

Zhu F, Leonard ED and Levin NW (2005): Body composition modelling in the calf using an equivalent circuit model of multi-frequency bioimpedance analysis. *Physiol Meas*, **26**:S133-S143.

Zhu F, Kuhlmann MK, Sarkar S (2004): Adjustment of dry weight in hemodialysis patients using intradialytic continuous multifrequency bioimpedance of the calf. *Journal of Artificial Organs*, **27**(2):104-109.

Zhu F, Hoenich NA, Kaysen G (2003): Measurement of intraperitoneal volume by segmental bioimpedance analysis during peritoneal dialysis. *Am J Kidney Diseases*, **42**:167-172.

Zhu F, Sarkar S, Kaitwatcharachai CH (2003): Methods and reproducibility of measurement of resistivity in the calf using regional bioimpedance analysis. *Blood Purif*, **21**:131-136.

Zhu, F Schneditz D, Kaufman AM and Levin NW (2000): Estimation of body fluid changes during peritoneal dialysis by segmental bioimpedance analysis. *Kidney Int*, **57**:299-306.

Zhu, F Schneditz D, Levin NW (1999): Sum of segmental bioimpedance analysis during ultrafiltration and hemodialysis reduces sensitivity to changes in body position. *Kidney Int*, **56** (2):692-699.

Zhu, F Schneditz D, Wang E, Martin K (1998): Validation of changes in extracellular volume measured during hemodialysis using a segmental bioimpedance technique. *ASAIO J*, M541-M545.

