

DEPARTAMENTO DE BIOLOGÍA CELULAR Y ANATOMÍA PATOLÓGICA
FACULTAD DE MEDICINA



*EL PAPEL DEL DIACILGLICEROL EN EL TRÁFICO DE MEMBRANAS EN LA ZONA
ENTRE EL RETÍCULO ENDOPLASMÁTICO Y EL COMPLEJO DE GOLGI*

Tesis presentada por Inés Fernández Ulibarri
para optar al título de Doctor por la Universidad de Barcelona

Bibliografía

1. A. T. Jones. (2007). Macropinocytosis: searching for an endocytic identity and role in the uptake of cell penetrating peptides. *J.Cell Mol.Med.* 11, 670-684
2. S. D. Conner and S. L. Schmid. (2003). Regulated portals of entry into the cell. *Nature* 422, 37-44
3. E. S. Seto, H. J. Bellen and T. E. Lloyd. (2002). When cell biology meets development: endocytic regulation of signaling pathways. *Genes Dev.* 16, 1314-1336
4. M. S. Robinson and J. S. Bonifacino. (2001). Adaptor-related proteins. *Curr.Opin.Cell Biol.* 13, 444-453
5. T. Kirchhausen. (2007). Making COPII coats. *Cell* 129, 1251-1252
6. F. M. Brodsky, C. Y. Chen, C. Knuehl and M. C. Towler, D. E. Wakeham. (2001). Biological basket weaving: formation and function of clathrin-coated vesicles. *Annu.Rev.Cell Dev.Biol.* 17, 517-568
7. B. van Deurs, K. Roepstorff, A. M. Hommelgaard and K. Sandvig. (2003). Caveolae: anchored, multifunctional platforms in the lipid ocean. *Trends Cell Biol.* 13, 92-100
8. J. B. Helms and C. Zurzolo. (2004). Lipids as targeting signals: lipid rafts and intracellular trafficking. *Traffic.* 5, 247-254
9. M. Edidin. (2001). Shrinking patches and slippery rafts: scales of domains in the plasma membrane. *Trends Cell Biol.* 11, 492-496
10. R. G. Anderson and K. Jacobson. (2002). A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains. *Science* 296, 1821-1825
11. F. R. Maxfield and T. E. McGraw. (2004). Endocytic recycling. *Nat.Rev.Mol.Cell Biol.* 5, 121-132
12. M. Kirkham and R. G. Parton (2005). Clathrin-independent endocytosis: new insights into caveolae and non-caveolar lipid raft carriers. *Biochim.Biophys.Acta* 1746, 349-363
13. R. B. Kelly. (1985). Pathways of protein secretion in eukaryotes. *Science* 230, 25-32
14. S. Ponnambalam and S. A. Baldwin. (2003). Constitutive protein secretion from the trans-Golgi network to the plasma membrane. *Mol.Membr.Biol.* 20, 129-139
15. M. Blazquez and K. I. Shennan. (2000). Basic mechanisms of secretion: sorting into the regulated secretory pathway. *Biochem.Cell Biol.* 78, 181-191
16. W. Nickel. (2005). Unconventional secretory routes: direct protein export across the plasma membrane of mammalian cells. *Traffic.* 6, 607-614
17. R. M. Rios and M. Bornens. (2003). The Golgi apparatus at the cell centre. *Curr.Opin.Cell Biol.* 15, 60-66
18. F. Kepes, A. Rambourg and B. Satiat-Jeunemaitre. (2005). Morphodynamics of the secretory pathway. *Int.Rev.Cytol.* 242, 55-120
19. G. Egea. (2001). Train trip into the cell: the Golgi Aparatus as the cetral station of the intracellular membrane traffic. *Ciencia al dia Internacional,* 4, 1-20

20. G. Egea and Rios RM. (2002). The role of the cytoskeleton in the structure and function of the Golgi apparatus. *The Golgi Apparatus- State of the art 110 years after Camillo Golgi's discovery*, 270-300 (Book chapter)
21. F. Lazaro-Dieguez. (2008). Implicación de los filamentos de actina en la arquitectura, homeostasis y tráfico de salida del aparato de Golgi y estudio de la formación y degradación de un agrosoma de actina (Thesis)
22. A. Hille-Rehfeld. (1995). Mannose 6-phosphate receptors in sorting and transport of lysosomal enzymes. *Biochim.Biophys.Acta* 1241, 177-194
23. J. H. Benting, A. G. Rietveld and K. Simons. (1999). N-Glycans mediate the apical sorting of a GPI-anchored, raft-associated protein in Madin-Darby canine kidney cells. *J.Cell Biol.* 146, 313-320
24. A. H. Merrill, Jr. (2002). De novo sphingolipid biosynthesis: a necessary, but dangerous, pathway. *J.Biol.Chem.* 277, 25843-25846
25. R. Schekman and L. Orci. (1996). Coat proteins and vesicle budding. *Science* 271, 1526-1533
26. C. Barlowe, *Biochim.* (1998). COPII and selective export from the endoplasmic reticulum. *Biophys.Acta* 1404, 67-76
27. L. Orci, M. Stamnes, M. Ravazzola, M. Amherdt, A. Perrelet, T.H. Sollner and J.E Rothman. (1997). Bidirectional transport by distinct populations of COPI-coated vesicles. *Cell* 90, 335-349
28. F. S. Gorelick and C. Shugrue (2001). Exiting the endoplasmic reticulum. *Mol.Cell Endocrinol.* 177, 13-18
29. A. Sesso, F. P. de Faria, E. S. Iwamura and H. Correa. (1994). A three-dimensional reconstruction study of the rough ER-Golgi interface in serial thin sections of the pancreatic acinar cell of the rat. *J.Cell Sci.* 107 (Pt 3), 517-528
30. G. Palade (1975). Intracellular aspects of the process of protein synthesis. *Science* 189, 347-358
31. C. Barlowe, L. Orci, T Yeung, M. Hosobuchi, S. Hamamoto, N. Salama, M.F Rexach, M. Ravazzola, M. Amherdt and R. Schekman. (1994). COPII: a membrane coat formed by Sec proteins that drive vesicle budding from the endoplasmic reticulum. *Cell* 77, 895-907
32. M. Aridor, J. Weissman, S. Bannykh, C. Nuoffer and W. E. Balch. (1998). Cargo selection by the COPII budding machinery during export from the ER. *J.Cell Biol.* 141, 61-70
33. S. I. Bannykh, N. Nishimura and W. E. Balch. (1998). Getting into the Golgi. *Trends Cell Biol.* 8, 21-25
34. H. P. Hauri and A. Schweizer. (1992). The endoplasmic reticulum-Golgi intermediate compartment. *Curr.Opin.Cell Biol.* 4, 600-608
35. C. Rabouille and J. Klumperman. (2005). Opinion: The maturing role of COPI vesicles in intra-Golgi transport. *Nat.Rev.Mol.Cell Biol.* 6, 812-817
36. S. Munro and H. R. Pelham. (1987). A C-terminal signal prevents secretion of luminal ER proteins. *Cell* 48, 899-907

37. J. C. Semenza, K. G. Hardwick, N. Dean and H. R. Pelham. (1990). ERD2, a yeast gene required for the receptor-mediated retrieval of luminal ER proteins from the secretory pathway. *Cell* 61, 1349-1357
38. H. R. Pelham. (1991). Recycling of proteins between the endoplasmic reticulum and Golgi complex. *Curr.Opin.Cell Biol.* 3, 585-591
39. C. Appenzeller, H. Andersson, F. Kappeler and H. P. Hauri. (1999). The lectin ERGIC-53 is a cargo transport receptor for glycoproteins. *Nat.Cell Biol.* 1, 330-334
40. M. C. Lee, E. A. Miller, J. Goldberg, L. Orci and R. Schekman. (2004). Bi-directional protein transport between the ER and Golgi. *Annu.Rev.Cell Dev.Biol.* 20, 87-123
41. L. Lippincott-Schwartz, J.G. Donaldson, A. Schweizer, E.G. Berger, H.P. Hauri, L.C. Yuan, and R.D. Klausner. (1990). Microtubule-dependent retrograde transport of proteins into the ER in the presence of brefeldin A suggests an ER recycling pathway. *Cell* 60, 821-836
42. B. Storrie, J. White, S. Rottger, E.H. Stelzer, T. Sukanuma and T. Nilsson. (1998). Recycling of golgi-resident glycosyltransferases through the ER reveals a novel pathway and provides an explanation for nocodazole-induced Golgi scattering. *J.Cell Biol.* 143, 1505-1521
43. B. Storrie. (2005). Maintenance of Golgi apparatus structure in the face of continuous protein recycling to the endoplasmic reticulum: making ends meet. *Int.Rev.Cytol.* 244, 69-94
44. R. B. Kelly. (1999). Deconstructing membrane traffic. *Trends Cell Biol.* 9, M29-M33
45. G. Warren and I. Mellman. (1999). Bulk flow redux?. *Cell* 98, 125-127
46. T. E. Kreis and R. Pepperkok. (1994). Coat proteins in intracellular membrane transport. *Curr.Opin.Cell Biol.* 6, 533-537
47. J. Bethune, M. Kol, J.Hoffmann, I. Reckmann, B. Brugger and F. Wieland. (2006). Coatomer, the coat protein of COPI transport vesicles, discriminates endoplasmic reticulum residents from p24 proteins. *Mol.Cell Biol.* 26, 8011-8021
48. J. Lippincott-Schwartz and W. Liu. (2006). Insights into COPI coat assembly and function in living cells. *Trends Cell Biol.* 16, e1-e4
49. L. Orci, M. Ravazzola, A. Volchuk, T. Engel, M. Gmachl, M. Amherdt, A. Perrelet, T.H. Sollner and J.E Rothman. (2000). Anterograde flow of cargo across the golgi stack potentially mediated via bidirectional "percolating" COPI vesicles. *Proc.Natl.Acad.Sci.U.S.A* 97, 10400-10405
50. M. J. Lewis and H. R. Pelham. (1990). A human homologue of the yeast HDEL receptor. *Nature* 348, 162-163
51. T. Nilsson, M. Jackson and P. A. Peterson. (1989). Short cytoplasmic sequences serve as retention signals for transmembrane proteins in the endoplasmic reticulum. *Cell* 58, 707-718
52. O. Martinez, C. Antony, G. Pehau-Arnaudet, E.G Berger, J. Salamero and B. Goud. (1997). GTP-bound forms of rab6 induce the redistribution of Golgi proteins into the endoplasmic reticulum. *Proc.Natl.Acad.Sci.U.S.A* 94, 1828-1833
53. A. Girod, B. Storrie, J.C. Simpson, L. Johannes, B. Goud, L.M Roberts, J.M Lord, T. Nilsson and R. Pepperkok. (1999). Evidence for a COP-I-independent transport route from the Golgi complex to the endoplasmic reticulum. *Nat.Cell Biol.* 1, 423-430

54. A. Echard, F. Jollivet, O. Martinez, J.J. Lacapere, A. Rousselet, I. Janoueix-Lerosey and B. Goud. (1998). Interaction of a Golgi-associated kinesin-like protein with Rab6. *Science* 279, 580-585
55. H. Miki, Y. Okada and N. Hirokawa. (2005). Analysis of the kinesin superfamily: insights into structure and function. *Trends Cell Biol.* 15, 467-476
56. T. Matanis, A. Akhmanova, P. Wulf, E. Del Nery, T. Weide, T. Stepanova, N. Galjart, F. Grosveld, B. Goud, C.I. De Zeeuw, A. Barnekow and C.C. Hoogenraad. (2002). Bicaudal-D regulates COPI-independent Golgi-ER transport by recruiting the dynein-dynactin motor complex. *Nat.Cell Biol.* 4, 986-992
57. A. Volchuk, M. Amherdt, M. Ravazzola, B. Brugger, V.M. Rivera, T. Clackson, A. Perrelet, T.H. Sollner, J.E. Rothman and L. Orci. (2000). Megavesicles implicated in the rapid transport of intracisternal aggregates across the Golgi stack. *Cell* 102, 335-348
58. H. R. Pelham and J. E. Rothman. (2000). The debate about transport in the Golgi--two sides of the same coin?. *Cell* 102, 713-719
59. L. Orci, B. S. Glick and J. E. Rothman. (1986). A new type of coated vesicular carrier that appears not to contain clathrin: its possible role in protein transport within the Golgi stack. *Cell* 46, 171-184
60. J. E. Rothman. (1996). The protein machinery of vesicle budding and fusion. *Protein Sci.* 5, 185-194
61. A. A. Mironov, P. Weidman and A. Luini. (1997). Variations on the intracellular transport theme: maturing cisternae and trafficking tubules. *J.Cell Biol.* 138, 481-484
62. A. Trucco, R.S. Polishchuk, O. Martella, A. Di Pentima, A. Fusella, D. Di Giandomenico E. San Pietro, G.V. Beznoussen E.V. Polishchuk, M. Baldassarre, R. Buccione, W.J. Geerts, A.J. Koster, K.N. Burger, A.A. Mironov and A. Luini. (2004). Secretory traffic triggers the formation of tubular continuities across Golgi sub-compartments. *Nat.Cell Biol.* 6, 1071-1081.
63. G. Vivero-Salmeron, J. Ballesta and J. A. Martinez-Menarguez(2008). Heterotypic tubular connections at the endoplasmic reticulum-Golgi complex interfac. *Histochem.Cell Biol.* 130, 709-717
64. M. A. De Matteis and A. Luini. (2008). Exiting the Golgi complex. *Nat.Rev.Mol.Cell Biol.* 9, 273-284
65. J. S. Bonifacino and L. M. Traub. (2003). Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annu.Rev.Biochem.* 72, 395-447
66. B. A. Potter, G. Ihrke, J. R. Bruns, K. M. Weixel and O. A. Weisz. (2004). Specific N-glycans direct apical delivery of transmembrane, but not soluble or glycosylphosphatidylinositol-anchored forms of endolyn in Madin-Darby canine kidney cells. *Mol.Biol.Cell* 15, 1407-1416
67. S. Schuck and K. Simons. (2004). Polarized sorting in epithelial cells: raft clustering and the biogenesis of the apical membrane. *J.Cell Sci.* 117, 5955-5964
68. P. Keller, D. Toomre, E. Diaz, J. White and K. Simons. (2001). Multicolour imaging of post-Golgi sorting and trafficking in live cells. *Nat.Cell Biol.* 3, 140-149
69. J. S. Bonifacino and J. Lippincott-Schwartz. (2003). Coat proteins: shaping membrane transport. *Nat.Rev.Mol.Cell Biol.* 4, 409-414

70. J. S. Bonifacino and B. S. Glick. (2004). The mechanisms of vesicle budding and fusion. *Cell* 116, 153-166
71. B.G Kearns, T.P. McGee, P. Mayinger, A. Gedvilaite, S.E. Phillips, S. Kagiwada and V.A. Bankaitis.(1997). Essential role for diacylglycerol in protein transport from the yeast Golgi complex. *Nature* 387, 101-105
72. M. S. Robinson. (2004). Adaptable adaptors for coated vesicles. *Trends Cell Biol.* 14, 167-174
73. J. E. Rothman. (2002). Lasker Basic Medical Research Award. The machinery and principles of vesicle transport in the cell. *Nat.Med.* 8, 1059-1062
74. J. E. Rothman. (1994). Mechanisms of intracellular protein transport. *Nature* 372, 55-63
75. R. Schekman and I. Mellman. (1997). Does COPI go both ways?. *Cell* 90, 197-200
76. W. Nickel, B. Brugger, F. T. Wieland. (1998). Protein and lipid sorting between the endoplasmic reticulum and the Golgi complex. *Semin.Cell Dev.Biol.* 9, 493-501
77. W. Nickel and F. T. Wieland. (1998). Biosynthetic protein transport through the early secretory pathway. *Histochem.Cell Biol.* 109, 477-486
78. W. E. Balch, R. A. Kahn and R. Schwaninger.(1992). ADP-ribosylation factor is required for vesicular trafficking between the endoplasmic reticulum and the cis-Golgi compartment. *J.Biol.Chem.* 267, 13053-13061
79. G. Tanigawa, L. Orci, M. Amherdt, M. Ravazzola, J.B. Helms and J.E Rothman (1993). Hydrolysis of bound GTP by ARF protein triggers uncoating of Golgi-derived COP-coated vesicles. *J.Cell Biol.* 123, 1365-1371
80. T. Kirchhausen. (2000). Three ways to make a vesicle.*Nat.Rev.Mol.Cell Biol.* 1, 187-198
81. J. Bigay, P. Gounon, S. Robineau and B. Antonny. (2003). Lipid packing sensed by ArfGAP1 couples COPI coat disassembly to membrane bilayer curvature. *Nature* 426, 563-566
82. C. Reinhard, M. Schweikert, F. T. Wieland and W. Nickel. (2003). Functional reconstitution of COPI coat assembly and disassembly using chemically defined components. *Proc.Natl.Acad.Sci.U.S.A* 100, 8253-8257
83. J. G. Donaldson, D. Finazzi and R. D. Klausner. (1992). Brefeldin A inhibits Golgi membrane-catalysed exchange of guanine nucleotide onto ARF protein. *Nature* 360, 350-352
84. J. B. Helms and J. E. Rothman. (1992). Inhibition by brefeldin A of a Golgi membrane enzyme that catalyses exchange of guanine nucleotide bound to ARF. *Nature* 360, 352-354
85. M. A. De Matteis and G. D'Angelo. (2007). The role of the phosphoinositides at the Golgi complex. *Biochem.Soc.Symp.* 74, 107-116
86. S. Lev. (2006). Lipid homeostasis and Golgi secretory function. *Biochem.Soc.Trans.* 34, 363-366
87. M. G. Roth. (1999). Lipid regulators of membrane traffic through the Golgi complex. *Trends Cell Biol.* 9, 174-179

88. C. D'Souza-Schorey and P. Chavrier. (2006). ARF proteins: roles in membrane traffic and beyond. *Nat.Rev.Mol.Cell Biol.* 7, 347-358
89. Z. Nie and P. A. Randazzo. (2006). Arf GAPs and membrane traffic. *J.Cell Sci.* 119, 1203-1211
90. H. T. McMahon and J. L. Gallop. (2005). Membrane curvature and mechanisms of dynamic cell membrane remodelling. *Nature* 438, 590-596
91. E. Rodriguez-Boulan, G. Kreitzer and A. Musch. (2005). Organization of vesicular trafficking in epithelia. *Nat.Rev.Mol.Cell Biol.* 6, 233-247
92. J. Lippincott-Schwartz, L. Yuan, C. Tipper, M. Amherdt, L. Orci and R.D. Klausner. (1991). Brefeldin A's effects on endosomes, lysosomes, and the TGN suggest a general mechanism for regulating organelle structure and membrane traffic. *Cell* 67, 601-616
93. H. Cao, S. Weller, J.D. Orth, J. Chen, B. Huang, J.L. Chen, M. Stamnes and M.A. McNiven. (2005). Actin and Arf1-dependent recruitment of a cortactin-dynamin complex to the Golgi regulates post-Golgi transport. *Nat.Cell Biol.* 7, 483-492
94. S. Carreno, A. E. Engqvist-Goldstein, C. X. Zhang, K. L. McDonald and D. G. Drubin. (2004). Actin dynamics coupled to clathrin-coated vesicle formation at the trans-Golgi network. *J.Cell Biol.* 165, 781-788
95. E. Kerkhoff, J.C. Simpson, C.B. Leberfinger, I.M. Otto, T. Doerks, P. Bork, U.R. Rapp, T. Raabe and R. Pepperkok. (2001). The Spir actin organizers are involved in vesicle transport processes. *Curr.Biol.* 11, 1963-1968
96. M. M. Kessels and B. Qualmann. (2004). The syndapin protein family: linking membrane trafficking with the cytoskeleton. *J.Cell Sci.* 117, 3077-3086
97. G. J. Praefcke and H. T. McMahon. (2004). The dynamin superfamily: universal membrane tubulation and fission molecules?. *Nat.Rev.Mol.Cell Biol.* 5, 133-147
98. A.L. Rozelle, L.M. Machesky, M. Yamamoto, M.H. Driessens, R.H. Insall, M.G. Roth, K. Luby-Phelps, G. Marriott, A. Hall and H.L. Yin. (2000). Phosphatidylinositol 4,5-bisphosphate induces actin-based movement of raft-enriched vesicles through WASP-Arp2/3. *Curr.Biol.* 10, 311-320
99. B. Antony. (2006). Membrane deformation by protein coats. *Curr.Opin.Cell Biol.* 18, 386-394
100. K. Farsad and P. De Camilli. (2003). Mechanisms of membrane deformation. *Curr.Opin.Cell Biol.* 15, 372-381
101. J.L. Gallop, C.C. Jao, H.M. Kent, P.J. Butler, P.R. Evans, R. Langen and H.T. McMahon. (2006). Mechanism of endophilin N-BAR domain-mediated membrane curvature. *EMBO J.* 25, 2898-2910
102. J. Lippincott-Schwartz and W. Liu. (2003). Membrane trafficking: coat control by curvature. *Nature* 426, 507-508
103. J. Zimmerberg and M. M. Kozlov. (2006). How proteins produce cellular membrane curvature. *Nat.Rev.Mol.Cell Biol.* 7, 9-19
104. E. Farge and P. F. Devaux. (1992). Shape changes of giant liposomes induced by an asymmetric transmembrane distribution of phospholipids. *Biophys.J.* 61, 347-357
105. K. N. Burger. (2000). Greasing membrane fusion and fission machineries. *Traffic.* 1, 605-613

106. D. Corda, C. C. Hidalgo, M. Bonazzi, A. Luini and S. Spano. (2002). Molecular aspects of membrane fission in the secretory pathway. *Cell Mol.Life Sci.* 59, 1819-1832
107. P. Helfrich and E. Jakobsson. (1990). Calculation of deformation energies and conformations in lipid membranes containing gramicidin channels. *Biophys.J.* 57, 1075-1084
108. M.A. De Matteis, M. Di Girolamo, A. Colanzi, M. Pallas, G. Di Tullio, L.J. McDonald, J. Moss, G. Santini, S. Bannykh and D. Corda. (1994). Stimulation of endogenous ADP-ribosylation by brefeldin A. *Proc.Natl.Acad.Sci.U.S.A* 91, 1114-1118
109. A. Mironov, A. Colanzi, M.G. Silletta, G. Fiucci, S. Flati, A. Fusella, R. Polishchuk, A.Jr. Mironov, G. Di Tullio, R. Weigert, V. Malhotra, D. Corda, M.A. De Matteis and A. Luini. (1997). Role of NAD⁺ and ADP-ribosylation in the maintenance of the Golgi structure. *J.Cell Biol.* 139, 1109-1118
110. R. Weigert, M.G. Silletta, S. Spano, G. Turacchio, C. Cericola, A. Colanzi, S. Senatore, R. Mancini, E.V. Polishchuk, M. Salmona, F. Facchiano, K.N. Burger, A. Mironov, A. Luini, and D. Corda. (1999). CtBP/BARS induces fission of Golgi membranes by acylating lysophosphatidic acid. *Nature* 402, 429-433
111. M. Bonazzi, S. Spano, G. Turacchio, C. Cericola, C. Valente, A. Colanzi, H.S. Kweon, V.W. Hsu, E.V. Polishchuck, R.S. Polishchuck, M. Sallese, T. Pulvirenti, D. Corda and A. Luini. (2005). CtBP3/BARS drives membrane fission in dynamin-independent transport pathways. *Nat.Cell Biol.* 7, 570-580
112. J.S. Yang, L. Zhang, S.Y. Lee, H. Gad, A. Luini and V.W. Hsu. (2006). Key components of the fission machinery are interchangeable. *Nat.Cell Biol.* 8, 1376-1382
113. J.s. Yang, S.Y. Lee, S. Spano, H. Gad, L. Zhang, Z. Nie, M. Bonazzi, D. Corda, A. Luini and V.W. Hsu. (2005). A role for BARS at the fission step of COPI vesicle formation from Golgi membrane. *EMBO J.* 24, 4133-4143
114. H. Inoue and P. A.Randazzo. (2007). Arf GAPs and their interacting proteins. *Traffic.* 8, 1465-1475
115. E. Cukierman, I. Huber, M. Rotman and D. Cassel. (1995). The ARF1 GTPase-activating protein: zinc finger motif and Golgi complex localization. *Science* 270, 1999-2002
116. J. Lanoix, J. Ouwendijk, A. Stark, E. Szafer, D. Cassel, K. Dejgaard, M. Weiss and T. Nilsson. (1999). Sorting of Golgi resident proteins into different subpopulations of COPI vesicles: a role for ArfGAP1. *EMBO J.* 18, 4935-4948
117. L. C. Bickford, E. Mossessova and J. Goldberg. (2004). A structural view of the COPII vesicle coat. *Curr.Opin.Struct.Biol.* 14, 147-153
118. W. Liu, R. Duden, R. D. Phair and J. Lippincott-Schwartz. (2005). ArfGAP1 dynamics and its role in COPI coat assembly on Golgi membranes of living cells. *J.Cell Biol.* 168, 1053-1063
119. T. Aoe, E. Cukierman, A. Lee, D. Cassel, P.J. Peters and V.W. Hsu. (1997). The KDEL receptor, ERD2, regulates intracellular traffic by recruiting a GTPase-activating protein for ARF1. *EMBO J.* 16, 7305-7316
120. T. Aoe, A. J. Lee, E. van Donselaar, P. J. Peters and V. W. Hsu. (1998). Modulation of intracellular transport by transported proteins: insight from regulation of COPI-mediated transport. *Proc.Natl.Acad.Sci.U.S.A* 95, 1624-1629

121. S. Y. Lee, J. S. Yang, W. Hong, R. T. Premont and V. W. Hsu. (2005). ARFGAP1 plays a central role in coupling COPI cargo sorting with vesicle formation. *J.Cell Biol.* 168, 281-290
122. P. J. Watson, G. Frigerio, B. M. Collins, R. Duden and D. J. Owen. (2004). Gamma-COP appendage domain - structure and function. *Traffic.* 5, 79-88
123. J. Goldberg. (1999). Structural and functional analysis of the ARF1-ARFGAP complex reveals a role for coatamer in GTP hydrolysis. *Cell* 96, 893-902
124. E. Szafer, M. Rotman and D. Cassel. (2001). Regulation of GTP hydrolysis on ADP-ribosylation factor-1 at the Golgi membrane. *J.Biol.Chem.* 276, 47834-47839
125. J. Bigay, J. F. Casella, G. Drin, B. Mesmin and B. Antonny. (2005). ArfGAP1 responds to membrane curvature through the folding of a lipid packing sensor motif. *EMBO J.* 24, 2244-2253
126. B. Mesmin, G. Drin, S. Levi, M. Rawet, D. Cassel, J. Bigay and B. Antonny. (2007). Two lipid-packing sensor motifs contribute to the sensitivity of ArfGAP1 to membrane curvature. *Biochemistry* 46, 1779-1790
127. A. Parnis, M. Rawet, L. Regev, B. Barkan, M. Rotman, M. Gaitner and D. Cassel. (2006). Golgi localization determinants in ArfGAP1 and in new tissue-specific ArfGAP1 isoforms. *J.Biol.Chem.* 281, 3785-3792
128. B. Antonny, I. Huber, S. Paris, M. Chabre and D. Cassel. (1997). Activation of ADP-ribosylation factor 1 GTPase-activating protein by phosphatidylcholine-derived diacylglycerols. *J.Biol.Chem.* 272, 30848-30851
129. Y.G. Chen, A. Siddhanta, C.D. Austin, S.M. Hammond, T.C. Sung, M.A. Frohman, A.J. Morris and D. Shield. (1997). Phospholipase D stimulates release of nascent secretory vesicles from the trans-Golgi network. *J.Cell Biol.* 138, 495-504
130. N. T. Ktistakis, H. A. Brown, M. G. Waters, P. C. Sternweis and M. G. Roth. (1996). Evidence that phospholipase D mediates ADP ribosylation factor-dependent formation of Golgi coated vesicles. *J.Cell Biol.* 134, 295-306
131. H. A. Brown, S. Gutowski, C. R. Moomaw, C. Slaughter and P. C. Sternweis. (1993). ADP-ribosylation factor, a small GTP-dependent regulatory protein, stimulates phospholipase D activity. *Cell* 75, 1137-1144
132. J. Stutchfield and S. Cockcroft. (1993). Correlation between secretion and phospholipase D activation in differentiated HL60 cells. *Biochem.J.* 293 (Pt 3), 649-655
133. T.R. Pettitt, A. Martin, T. Horton, C. Liossis, J.M. Lord and M.J. Wakelam. (1997). Diacylglycerol and phosphatidate generated by phospholipases C and D, respectively, have distinct fatty acid compositions and functions. Phospholipase D-derived diacylglycerol does not activate protein kinase C in porcine aortic endothelial cells. *J.Biol.Chem.* 272, 17354-17359
134. D. D. Sabatini, M. Adesnik, I. E. Ivanov and J. P. Simon. (1996). Mechanism of formation of post Golgi vesicles from TGN membranes: Arf-dependent coat assembly and PKC-regulated vesicle scission. *Biocell* 20, 287-300
135. J. P. Simon, I. E. Ivanov, M. Adesnik and D. D. Sabatini. (1996). The production of post-Golgi vesicles requires a protein kinase C-like molecule, but not its phosphorylating activity. *J.Cell Biol.* 135, 355-370
136. J.P. Simon, I.E. Ivanov, B. Shopsin, D. Hersh, M. Adesnik and D.D. Sabatini. (1996). The in vitro generation of post-Golgi vesicles carrying viral envelope glycoproteins

- requires an ARF-like GTP-binding protein and a protein kinase C associated with the Golgi apparatus. *J.Biol.Chem.* 271, 16952-16961
137. K. M. Conricode, K. A. Brewer and J. H. Exton. (1992). Activation of phospholipase D by protein kinase C. Evidence for a phosphorylation-independent mechanism. *J.Biol.Chem.* 267, 7199-7202
 138. W. D. Singer, H. A. Brown, X. Jiang and P. C. Sternweis. (1996). Regulation of phospholipase D by protein kinase C is synergistic with ADP-ribosylation factor and independent of protein kinase activity. *J.Biol.Chem.* 271, 4504-4510
 139. M. Liljedahl, Y. Maeda, A. Colanzi, I. Ayala, J. Van Lint and V. Malhotra. (2001). Protein kinase D regulates the fission of cell surface destined transport carriers from the trans-Golgi network. *Cell* 104, 409-420
 140. J. Prestle, K. Pfizenmaier, J. Brenner and F. J. Johannes. (1996). Protein kinase C mu is located at the Golgi compartment. *J.Cell Biol.* 134, 1401-1410
 141. J. Van Lint, A. Rykx, Y. Maeda, T. Vantus, S. Sturany, V. Malhotra, J.R. Vandenheede and T. Seufferlein. (2002). Protein kinase D: an intracellular traffic regulator on the move. *Trends Cell Biol.* 12, 193-200
 142. J. V. Van Lint, J. Sinnett-Smith and E. Rozengurt. (1995). Expression and characterization of PKD, a phorbol ester and diacylglycerol-stimulated serine protein kinase. *J.Biol.Chem.* 270, 1455-1461
 143. C. L. Baron and V. Malhotra. (2002). Role of diacylglycerol in PKD recruitment to the TGN and protein transport to the plasma membrane. *Science* 295, 325-328
 144. Y. Maeda, G. V. Beznoussenko, J. Van Lint, A. A. Mironov and V. Malhotra (2001). Recruitment of protein kinase D to the trans-Golgi network via the first cysteine-rich domain. *EMBO J.* 20, 5982-5990
 145. W. B. Huttner and A. Schmidt. (2000). Lipids, lipid modification and lipid-protein interaction in membrane budding and fission--insights from the roles of endophilin A1 and synaptophysin in synaptic vesicle endocytosis. *Curr.Opin.Neurobiol.* 10, 543-551
 146. W. Dowhan. (1997). Molecular basis for membrane phospholipid diversity: why are there so many lipids?. *Annu.Rev.Biochem.* 66, 199-232
 147. T. de Vrije, R. L. de Swart, W. Dowhan, J. Tommassen and B. de Kruijff. (1988). Phosphatidylglycerol is involved in protein translocation across Escherichia coli inner membranes. *Nature* 334, 173-175
 148. W. van Klompenburg, I. Nilsson, G. von Heijne and B. de Kruijff. (1997). Anionic phospholipids are determinants of membrane protein topolog. *EMBO J.* 16, 4261-4266
 149. S. H. White, A. S. Ladokhin, S. Jayasinghe and K. Hristova. (2001). How membranes shape protein structure. *J.Biol.Chem.* 276, 32395-32398
 150. L. Heginbotham, L. Kolmakova-Partensky and C. Miller. (1998). Functional reconstitution of a prokaryotic K⁺ channel. *J.Gen.Physiol* 111, 741-749
 151. F. I. Valiyaveetil, Y. Zhou and R. MacKinnon. (2002). Lipids in the structure, folding, and function of the KcsA K⁺ channel. *Biochemistry* 41, 10771-10777
 152. Y. Zhou, J. H. Morais-Cabral, A. Kaufman and R. MacKinnon. (2001). Chemistry of ion coordination and hydration revealed by a K⁺ channel-Fab complex at 2.0 Å resolution. *Nature* 414, 43-48

153. A. G. Lee. (2004). How lipids affect the activities of integral membrane proteins. *Biochim.Biophys.Acta* 1666, 62-87
154. van den Brink-van der Laan, J. A. Killian and B. de Kruijff. (2004). Nonbilayer lipids affect peripheral and integral membrane proteins via changes in the lateral pressure profile. *Biochim.Biophys.Acta* 1666, 275-288
155. S. Cockcroft. (2001). Signalling roles of mammalian phospholipase D1 and D2. *Cell Mol.Life Sci.* 58, 1674-1687
156. T. Munnik, R. F. Irvine and A. Musgrave. (1998). Phospholipid signalling in plants. *Biochim.Biophys.Acta* 1389, 222-272
157. M. Manifava, J.W. Thuring, Z.Y. Lim, L. Packman, A.B. Holmes and N.T. Ktistakis. (2001). Differential binding of traffic-related proteins to phosphatidic acid- or phosphatidylinositol (4,5)- bisphosphate-coupled affinity reagents. *J.Biol.Chem.* 276, 8987-8994
158. J. C. Holthuis and T. P. Levine. (2005). Lipid traffic: floppy drives and a superhighway. *Nat.Rev.Mol.Cell Biol.* 6, 209-220
159. G. van Meer, D. R. Voelker and G. W. Feigenson. (2008). Membrane lipids: where they are and how they behave. *Nat.Rev.Mol.Cell Biol.* 9, 112-124
160. D. Voet and Voet J.G. (1995). *Biochemistry*, I. John Wiley&Sons, Ed.
161. E.M. Deacon, T.R. Pettitt, P. Webb, T. Cross, H. Chahal, M.J. Wakelam and J.M. Lord. (2002). Generation of diacylglycerol molecular species through the cell cycle: a role for 1-stearoyl, 2-arachidonyl glycerol in the activation of nuclear protein kinase C-betaII at G2/M. *J.Cell Sci.* 115, 983-989
162. S. Ichikawa and Y. Hirabayashi. (1998). Glucosylceramide synthase and glycosphingolipid synthesis. *Trends Cell Biol.* 8, 198-202
163. D. N. Brindley and D. W. Waggoner. (1998). Mammalian lipid phosphate phosphohydrolases. *J.Biol.Chem.* 273, 24281-24284
164. S. Pyne, K. C. Kong and P. I. Darroch. (2004). Lysophosphatidic acid and sphingosine 1-phosphate biology: the role of lipid phosphate phosphatases *Semin.Cell Dev.Biol.* 15, 491-501
165. S. G. Rhee. (2001). Regulation of phosphoinositide-specific phospholipase C. *Annu.Rev.Biochem.* 70, 281-312
166. A. K. Thompson, S. P. Mostafapour, L. C. Denlinger, J. E. Bleasdale and S. K. Fisher. (1991). The aminosteroid U-73122 inhibits muscarinic receptor sequestration and phosphoinositide hydrolysis in SK-N-SH neuroblastoma cells. A role for Gp in receptor compartmentation. *J.Biol.Chem.* 266, 23856-23862
167. S. Carrasco and I. Merida. (2004). Diacylglycerol-dependent binding recruits PKCtheta and RasGRP1 C1 domains to specific subcellular localizations in living T lymphocyte. *Mol.Biol.Cell* 15, 2932-2942
168. R. E. Pagano. (1988). What is the fate of diacylglycerol produced at the Golgi apparatus?. *Trends Biochem.Sci.* 13, 202-205
169. V. A. Bankaitis. (2002). Cell biology. Slick recruitment to the Golgi. *Science* 295, 290-291

170. V. Litvak, N. Dahan, S. Ramachandran, H. Sabanay and S. Lev. (2005). Maintenance of the diacylglycerol level in the Golgi apparatus by the Nir2 protein is critical for Golgi secretory function. *Nat.Cell Biol.* 7, 225-234
171. I. Fernandez-Ulibarri, M. Vilella, F. Lazaro-Diequez, E. Sarri, S.E. Martinez, N. Jimenez, E. Claro, I. Merida, K.N. Burger and G. Egea. (2007). Diacylglycerol is required for the formation of COPI vesicles in the Golgi-to-ER transport pathway. *Mol.Biol.Cell* 18, 3250-3263
172. J. H. Exton. (1999). Regulation of phospholipase D. *Biochim.Biophys.Acta* 1439, 121-133
173. G. M. Jenkins and M. A. Frohman. (2005). Phospholipase D: a lipid centric review. *Cell Mol.Life Sci.* 62, 2305-2316
174. D. Jones, C. Morgan and S. Cockcroft. (1999). Phospholipase D and membrane traffic. Potential roles in regulated exocytosis, membrane delivery and vesicle budding. *Biochim.Biophys.Acta* 1439, 229-244
175. G. Du, Y.M. Altshuller, N. Vitale, P. Huang, S. Chasserot-Golaz, A.J. Morris, M.F. Bader and M.A. Frohman. (2003). Regulation of phospholipase D1 subcellular cycling through coordination of multiple membrane association motifs. *J.Cell Biol.* 162, 305-315
176. V.A. Sciorra, S.A. Rudge, J. Wang, S. McLaughlin, J. Engebrecht and A.J. Morris. (2002). Dual role for phosphoinositides in regulation of yeast and mammalian phospholipase D enzymes. *J.Cell Biol.* 159, 1039-1049
177. J. M. Sugars, S. Celtek, M. Manifava, J. Coadwell and N. T. Ktistakis. (2002). Hierarchy of membrane-targeting signals of phospholipase D1 involving lipid modification of a pleckstrin homology domain. *J.Biol.Chem.* 277, 29152-29161
178. M. A. Frohman, T. C. Sung and A. J. Morris. (1999). Mammalian phospholipase D structure and regulation. *Biochim.Biophys.Acta* 1439, 175-186
179. M. Liscovitch, M. Czarny, G. Fiucci, Y. Lavie and X. Tang. (1999). Localization and possible functions of phospholipase D isozymes. *Biochim.Biophys.Acta* 1439, 245-263
180. M. McDermott, M. J. Wakelam and A. J. Morris. (2004). Phospholipase D. *Biochem.Cell Biol.* 82, 225-253
181. M. Manifava, J. Sugars and N. T. Ktistakis. (1999). Modification of catalytically active phospholipase D1 with fatty acid in vivo. *J.Biol.Chem.* 274, 1072-1077
182. Z. Freyberg, D. Sweeney, A. Siddhanta, S. Bourgoin, M. Frohman and D. Shields.(2001). Intracellular localization of phospholipase D1 in mammalian cells. *Mol.Biol.Cell* 12, 943-955
183. Z. Freyberg, S. Bourgoin and D. Shields. (2002). Phospholipase D2 is localized to the rims of the Golgi apparatus in mammalian cells. *Mol.Biol.Cell* 13, 3930-3942
184. D. N. Brindley. (1984). Intracellular translocation of phosphatidate phosphohydrolase and its possible role in the control of glycerolipid synthesis. *Prog.Lipid Res.* 23, 115-133
185. G. M. Carman and S. A. Henry. (1999). Phospholipid biosynthesis in the yeast *Saccharomyces cerevisiae* and interrelationship with other metabolic processes. *Prog.Lipid Res.* 38, 361-399

186. D. Sorger and G. Daum. (2003). Triacylglycerol biosynthesis in yeast. *Appl.Microbiol.Biotechnol.* 61, 289-299
187. D. N. Brindley. (2004). Lipid phosphate phosphatases and related proteins: signaling functions in development, cell division, and cancer. *J.Cell Biochem.* 92, 900-912
188. Z. Jamal, A. Martin, A. Gomez-Munoz and D. N. Brindley. (1991). Plasma membrane fractions from rat liver contain a phosphatidate phosphohydrolase distinct from that in the endoplasmic reticulum and cytosol. *J.Biol.Chem.* 266, 2988-2996
189. A. Gomez-Munoz, E. H. Hamza and D. N. Brindley. (1992). Effects of sphingosine, albumin and unsaturated fatty acids on the activation and translocation of phosphatidate phosphohydrolases in rat hepatocytes. *Biochim.Biophys.Acta* 1127, 49-56
190. A. Gomez-Munoz, G.M. Hatch, A. Martin, Z. Jamal, D.E. Vance and D.N. Brindley.(1992). Effects of okadaic acid on the activities of two distinct phosphatidate phosphohydrolases in rat hepatocytes. *FEBS Lett.* 301, 103-106
191. O. Aridor-Piterman, Y. Lavie and M. Liscovitch. (1992). Bimodal distribution of phosphatidic acid phosphohydrolase in NG108-15 cells. Modulation by the amphiphilic lipids oleic acid and sphingosine. *Eur.J.Biochem.* 204, 561-568
192. J. S. Long, N. J. Pyne and S. Pyne. (2008). Lipid phosphate phosphatases form homo- and hetero-oligomers: catalytic competency, subcellular distribution and function. *Biochem.J.* 411, 371-377
193. I. N. Fleming and S. J. Yeaman. (1995). Purification and characterization of N-ethylmaleimide-insensitive phosphatidic acid phosphohydrolase (PAP2) from rat liver. *Biochem.J.* 308 (Pt 3), 983-989
194. H. Kanoh, S. Imai, K. Yamada and F. Sakane. (1992). Purification and properties of phosphatidic acid phosphatase from porcine thymus membranes. *J.Biol.Chem.* 267, 25309-25314
195. D. W. Waggoner, A. Martin, J. Dewald, A. Gomez-Munoz and D. N. Brindley. (1995). Purification and characterization of novel plasma membrane phosphatidate phosphohydrolase from rat liver. *J.Biol.Chem.* 270, 19422-19429
196. D. K. Perry, V. L. Stevens, T. S. Widlanski and J. D. Lambeth. (1993). A novel ecto-phosphatidic acid phosphohydrolase activity mediates activation of neutrophil superoxide generation by exogenous phosphatidic acid. *J.Biol.Chem.* 268, 25302-25310
197. D. W. Waggoner, A. Gomez-Munoz, J. Dewald and D. N. Brindley. (1996). Phosphatidate phosphohydrolase catalyzes the hydrolysis of ceramide 1-phosphate, lysophosphatidate, and sphingosine 1-phosphate. *J.Biol.Chem.* 271, 16506-16509
198. M. Xie and M. G. Low. (1994). Identification and characterization of an ecto-(lyso) phosphatidic acid phosphatase in PAM212 keratinocytes. *Arch.Biochem.Biophys.* 312, 254-259
199. D. Barila, M. Plateroti, F. Nobili, A.O. Muda, Y. Xie, T. Morimoto and G. Perozzi. (1996). The Dri 42 gene, whose expression is up-regulated during epithelial differentiation, encodes a novel endoplasmic reticulum resident transmembrane protein. *J.Biol.Chem.* 271, 29928-29936
200. Q. X. Zhang, C. S. Pilquill, J. Dewald, L. G. Berthiaume and D. N. Brindley. (2000). Identification of structurally important domains of lipid phosphate phosphatase-1: implications for its sites of action. *Biochem.J.* 345 Pt 2, 181-184

201. R. J. Tate, D. Tolan and S. Pyne. (1999). Molecular cloning of magnesium-independent type 2 phosphatidic acid phosphatases from airway smooth muscle. *Cell Signal.* 11, 515-522
202. R. Roberts, V. A. Sciorra and A. J. Morris. (1998). Human type 2 phosphatidic acid phosphohydrolases. Substrate specificity of the type 2a, 2b, and 2c enzymes and cell surface activity of the 2a isoform. *J.Biol.Chem.* 273, 22059-22067
203. M. Kai, I. Wada, S. Imai, F. Sakane and H. Kanoh. (1996). Identification and cDNA cloning of 35-kDa phosphatidic acid phosphatase (type 2) bound to plasma membranes. Polymerase chain reaction amplification of mouse H₂O₂-inducible hic53 clone yielded the cDNA encoding phosphatidic acid phosphatase. *J.Biol.Chem.* 271, 18931-18938
204. M. Kai, I. Wada, S. Imai, F. Sakane and H. Kanoh. (1997). Cloning and characterization of two human isozymes of Mg²⁺-independent phosphatidic acid phosphatase. *J.Biol.Chem.* 272, 24572-24578
205. V. A. Sciorra and A. J. Morris. (1999). Sequential actions of phospholipase D and phosphatidic acid phosphohydrolase 2b generate diglyceride in mammalian cells. *Mol.Biol.Cell* 10, 3863-3876
206. F. Alderton, P. Darroch, B. Sambhi, A. McKie, I.S. Ahmed, N. Pyne and S. Pyne. (2001). G-protein-coupled receptor stimulation of the p42/p44 mitogen-activated protein kinase pathway is attenuated by lipid phosphate phosphatases 1, 1a, and 2 in human embryonic kidney 293 cells. *J.Biol.Chem.* 276, 13452-13460
207. M. Nanjundan and F. Possmayer. (2001). Pulmonary lipid phosphate phosphohydrolase in plasma membrane signalling platforms. *Biochem.J.* 358, 637-646
208. J. Long, P. Darroch, K.F. Wan, K.C. Kong, N. Ktistakis, N.J. Pyne, S. Pyne. (2005). Regulation of cell survival by lipid phosphate phosphatases involves the modulation of intracellular phosphatidic acid and sphingosine 1-phosphate pools. *Biochem.J.* 391, 25-32
209. R. Jasinska, Q.X. Zhang, C. Pilquill, I. Singh, J. Xu, J. Dewald, D.A. Dillon, L.G. Berthiaume, G.M. Carman, D.W. Waggoner and D.N. Brindley. (1999). Lipid phosphate phosphohydrolase-1 degrades exogenous glycerolipid and sphingolipid phosphate esters. *Biochem.J.* 340 (Pt 3), 677-686
210. T. Ishikawa, M. Kai, I. Wada and H. Kanoh. (2000). Cell surface activities of the human type 2b phosphatidic acid phosphatase. *J.Biochem.* 127, 645-651
211. Y. J. Jia, M. Kai, I. Wada, F. Sakane and H. Kanoh. (2003). Differential localization of lipid phosphate phosphatases 1 and 3 to cell surface subdomains in polarized MDCK cells. *FEBS Lett.* 552, 240-246
212. S.S. Smyth, V.A. Sciorra, Y.J. Sigal, Z. Pamuklar, Z. Wang, Y. Xu, G.D. Prestwich and A.J. Morris (2003). Lipid phosphate phosphatases regulate lysophosphatidic acid production and signaling in platelets: studies using chemical inhibitors of lipid phosphate phosphatase activity. *J.Biol.Chem.* 278, 43214-43223
213. C. Burnett, P. Makridou, L. Hewlett and K. Howard. (2004). Lipid phosphate phosphatases dimerise, but this interaction is not required for in vivo activity. *BMC.Biochem.* 5, 2
214. A. Martin, A. Gomez-Munoz, D. W. Waggoner, J. C. Stone and D. N. Brindley. (1993). Decreased activities of phosphatidate phosphohydrolase and phospholipase D in ras and tyrosine kinase (fps) transformed fibroblasts. *J.Biol.Chem.* 268, 23924-23932

215. F. M. Goni and A. Alonso. (1999). Structure and functional properties of diacylglycerols in membranes. *Prog.Lipid Res.* 38, 1-48
216. J. H. Exton. (1994). Phosphatidylcholine breakdown and signal transduction. *Biochim.Biophys.Acta* 1212, 26-42
217. S. A. Barker, K. K. Caldwell, J. R. Pfeiffer and B. S. Wilson. (1998). Wortmannin-sensitive phosphorylation, translocation, and activation of PLCgamma1, but not PLCgamma2, in antigen-stimulated RBL-2H3 mast cells. *Mol.Biol.Cell* 9, 483-496
218. T.G. Jin, T. Satoh, Y. Liao, C. Song, X. Gao, K. Kariya, C.D. Hu and T. Kataoka. (2001). Role of the CDC25 homology domain of phospholipase Cepsilon in amplification of Rap1-dependent signaling. *J.Biol.Chem.* 276, 30301-30307
219. M. Ohashi, de V. Jan, R. Frank, G. Snoek, V. Bankaitis, K. Wirtz and W.B. Huttner. (1995). A role for phosphatidylinositol transfer protein in secretory vesicle formation. *Nature* 377, 544-547
220. B. G. Kearns, J. G. Alb, Jr. and V. Bankaitis. (1998). Phosphatidylinositol transfer proteins: the long and winding road to physiological function. *Trends Cell Biol.* 8, 276-282
221. X. Li, Z. Xie and V. A. Bankaitis. (2000). Phosphatidylinositol / phosphatidylcholine transfer proteins in yeast. *Biochim.Biophys.Acta* 1486, 55-71
222. C. Wiedemann and S. Cockcroft. (1998). The Role of Phosphatidylinositol Transfer Proteins (PITPs) in Intracellular Signalling. *Trends Endocrinol.Metab* 9, 324-328
223. C. Ramoni, F. Spadaro, M. Menegon and F. Podo. (2001). Cellular localization and functional role of phosphatidylcholine-specific phospholipase C in NK cells. *J.Immunol.* 167, 2642-2650
224. K. Huitema, D. J. van den, J. F. Brouwers and J. C. Holthuis. (2004). Identification of a family of animal sphingomyelin synthases. *EMBO J.* 23, 33-44
225. S. Yamaoka, M. Miyaji, T. Kitano, H. Umehara and T. Okazaki. (2004). Expression cloning of a human cDNA restoring sphingomyelin synthesis and cell growth in sphingomyelin synthase-defective lymphoid cells. *J.Biol.Chem.* 279, 18688-18693
226. M. Villani, M. Subathra, Y.B. Im, Y. Choi, P. Signorelli, M. Del Poeta, C. Luberto. (2008). Sphingomyelin synthases regulate production of diacylglycerol at the Golgi. *Biochem.J.* 414, 31-41
227. Y. Ono, T. Fujii, K. Igarashi, T. Kuno, C. Tanaka, U. Kikkawa and Y. Nishizuka. (1989). Phorbol ester binding to protein kinase C requires a cysteine-rich zinc-finger-like sequence. *Proc.Natl.Acad.Sci.U.S.A* 86, 4868-4871
228. D. J. Burns and R. M. Bell. (1991). Protein kinase C contains two phorbol ester binding domains. *J.Biol.Chem.* 266, 18330-18338
229. A. F. Quest, E. S. Bardes and R. M. Bell. (1994). A phorbol ester binding domain of protein kinase C gamma. High affinity binding to a glutathione-S-transferase/Cys2 fusion protein. *J.Biol.Chem.* 269, 2953-2960
230. M. J. Caloca, J. L. Zugaza and X. R. Bustelo. (2003). Exchange factors of the RasGRP family mediate Ras activation in the Golgi. *J.Biol.Chem.* 278, 33465-33473
231. M. J. Caloca, H. Wang, A. Delemos, S. Wang and M. G. Kazanietz. (2001). Phorbol esters and related analogs regulate the subcellular localization of beta 2-chimaerin, a non-protein kinase C phorbol ester receptor. *J.Biol.Chem.* 276, 18303-18312

232. H. Wang and M. G. Kazanietz. (2002). Chimaerins, novel non-protein kinase C phorbol ester receptors, associate with Tmp21-I (p23): evidence for a novel anchoring mechanism involving the chimaerin C1 domain. *J.Biol.Chem.* 277, 4541-4550
233. T. Kajimoto, Y. Shirai, N. Sakai, T. Yamamoto, H. Matsuzaki, U. Kikkawa and N. Saito. (2004). Ceramide-induced apoptosis by translocation, phosphorylation, and activation of protein kinase Cdelta in the Golgi complex. *J.Biol.Chem.* 279, 12668-12676
234. B. Sparatore, M. Passalacqua, M. Pedrazzi, S. Ledda, M. Patrone, D. Gaggero, S. Pontremoli and E. Melloni. (2003). Role of the kinase activation loop on protein kinase C theta activity and intracellular localisation. *FEBS Lett.* 554, 35-40
235. R.V. Stahelin, M.A. Digman, M. Medkova, B. Ananthanarayanan, H.R. Melowic, J.D. Rafter and W. Cho.(2005). Diacylglycerol-induced membrane targeting and activation of protein kinase Cepsilon: mechanistic differences between protein kinases Cdelta and Cepsilon. *J.Biol.Chem.* 280, 19784-19793
236. C. Lehel, Z. Olah, G. Jakab and W. B. Anderson. (1995). Protein kinase C epsilon is localized to the Golgi via its zinc-finger domain and modulates Golgi function. *Proc.Natl.Acad.Sci.U.S.A* 92, 1406-1410
237. A. Schultz, M. Ling and C. Larsson. (2004). Identification of an amino acid residue in the protein kinase C C1b domain crucial for its localization to the Golgi network. *J.Biol.Chem.* 279, 31750-31760
238. L. M. Keranen, E. M. Dutil and A. C. Newton. (1995). Protein kinase C is regulated in vivo by three functionally distinct phosphorylations. *Curr.Biol.* 5, 1394-1403
239. T. R. Xu, G. He, K. Dobson, K. England and M. Rumsby. (2007). Phosphorylation at Ser729 specifies a Golgi localisation for protein kinase C epsilon (PKCepsilon) in 3T3 fibroblasts. *Cell Signal.* 19, 1986-1995
240. D. B. Parekh, W. Ziegler and P. J. Parker. (2000). Multiple pathways control protein kinase C phosphorylation. *EMBO J.* 19, 496-503
241. C. Lehel, Z. Olah, G. Petrovics, G. Jakab and W. B. Anderson. (1996). Influence of various domains of protein kinase C epsilon on its PMA-induced translocation from the Golgi to the plasma membrane. *Biochem.Biophys.Res.Commun.* 223, 98-103
242. K. England and M. G. Rumsby. (2000). Changes in protein kinase C epsilon phosphorylation status and intracellular localization as 3T3 and 3T6 fibroblasts grow to confluency and quiescence: a role for phosphorylation at ser-729?. *Biochem.J.* 352 Pt 1, 19-26
243. D. Ron and D. Mochly-Rosen. (1994). Agonists and antagonists of protein kinase C function, derived from its binding proteins. *J.Biol.Chem.* 269, 21395-21398
244. D. Ron, C.H. Chen, J. Caldwell, L. Jamieson, E. Orr and D. Mochly-Rosen. (1994). Cloning of an intracellular receptor for protein kinase C: a homolog of the beta subunit of G proteins. *Proc.Natl.Acad.Sci.U.S.A* 91, 839-843
245. M. Csukai, C. H. Chen, M. A. De Matteis and D. Mochly-Rosen. (1997). The coatomer protein beta'-COP, a selective binding protein (RACK) for protein kinase Cepsilon. *J.Biol.Chem.* 272, 29200-29206
246. T. Shemesh, A. Luini, V. Malhotra, K. N. Burger and M. M. Kozlov. (2003). Prefission constriction of Golgi tubular carriers driven by local lipid metabolism: a theoretical model. *Biophys.J.* 85, 3813-3827

247. F. Bard and V. Malhotra. (2006). The formation of TGN-to-plasma-membrane transport carriers. *Annu.Rev.Cell Dev.Biol.* 22, 439-455
248. S. L. Hofmann and P. W. Majerus. (1982). Modulation of phosphatidylinositol-specific phospholipase C activity by phospholipid interactions, diglycerides, and calcium ions. *J.Biol.Chem.* 257, 14359-14364
249. A. V. Villar, F. M. Goni and A. Alonso. (2001). Diacylglycerol effects on phosphatidylinositol-specific phospholipase C activity and vesicle fusion. *FEBS Lett.* 494, 117-120
250. R. M. Kramer, G. C. Checani and D. Deykin. (1987). Stimulation of Ca²⁺-activated human platelet phospholipase A2 by diacylglycerol. *Biochem.J.* 248, 779-783
251. R. M. Epand and D. S. Lester. (1990). The role of membrane biophysical properties in the regulation of protein kinase C activity. *Trends Pharmacol.Sci.* 11, 317-320
252. S.J. Slater, M.B. Kelly, F.J. Taddeo, C. Ho, E. Rubin and C.D. Stubbs. (1994). The modulation of protein kinase C activity by membrane lipid bilayer structure. *J.Biol.Chem.* 269, 4866-4871
253. B. E. Slack, J. Brey and R. J. Wurtman. (1991). Production of diacylglycerol by exogenous phospholipase C stimulates CTP:phosphocholine cytidyltransferase activity and phosphatidylcholine synthesis in human neuroblastoma cells. *J.Biol.Chem.* 266, 24503-24508
254. G. S. Attard, R. H. Templer, W. S. Smith, A. N. Hunt and S. Jackowski. (2000). Modulation of CTP:phosphocholine cytidyltransferase by membrane curvature elastic stress. *Proc.Natl.Acad.Sci.U.S.A* 97, 9032-9036
255. R. S. Arnold, A. A. DePaoli-Roach and R. B. Cornell. (1997). Binding of CTP:phosphocholine cytidyltransferase to lipid vesicles: diacylglycerol and enzyme dephosphorylation increase the affinity for negatively charged membranes. *Biochemistry* 36, 6149-6156
256. E. E. Kooijman, V. Chupin, B. de Kruijff and K. N. Burger. (2003). Modulation of membrane curvature by phosphatidic acid and lysophosphatidic acid. *Traffic.* 4, 162-174
257. E.E. Kooijman, V. Chupin, N.L Fuller, M.M. Kozlov, B. de Kruijff, K.N. Burger and P.R. Rand. (2005). Spontaneous curvature of phosphatidic acid and lysophosphatidic acid. *Biochemistry* 44, 2097-2102
258. A. Schmidt, M. Wolde, C. Thiele, W. Fest, H. Kratzin, A.V. Podtelejnikov, W. Witke, W.B Huttner and H.D. Soling (1999). Endophilin I mediates synaptic vesicle formation by transfer of arachidonate to lysophosphatidic acid. *Nature* 401, 133-141
259. K. Bi, M. G. Roth and N. T. Ktistakis. (1997). Phosphatidic acid formation by phospholipase D is required for transport from the endoplasmic reticulum to the Golgi complex. *Curr.Biol.* 7, 301-307
260. C.C. Hidalgo, M. Bonazzi, S. Spano, G.Turacchio, A. Colanzi, A. Luini, and D. Corda. (2004). Mitotic Golgi partitioning is driven by the membrane-fissioning protein CtBP3/BARS. *Science* 305, 93-96
261. Y. Jun, R. A. Fratti and W. Wickner. (2004). Diacylglycerol and its formation by phospholipase C regulate Rab- and SNARE-dependent yeast vacuole fusion. *J.Biol.Chem.* 279, 53186-53195
262. J.L. Nieva, A. Alonso, G. Basanez, F.M. Goni, A. Gulik, R. Vargas and V. Luzzati. (1995). Topological properties of two cubic phases of a

- phospholipid:cholesterol:diacylglycerol aqueous system and their possible implications in the phospholipase C-induced liposome fusion. *FEBS Lett.* 368, 143-147
263. M. B. Ruiz-Arguello, G. Basanez, F. M. Goni and A. Alonso. (1996). Different effects of enzyme-generated ceramides and diacylglycerols in phospholipid membrane fusion and leakage. *J.Biol.Chem.* 271, 26616-26621
264. A. Siddhanta and D. Shields. (1998). Secretory vesicle budding from the trans-Golgi network is mediated by phosphatidic acid levels. *J.Biol.Chem.* 273, 17995-17998
265. R. Weigert, M.G. Silletta, S. Spano, G. Turacchio, C. Cericola, A. Colanzi, S. Senatore, R. Mancini, E.V. Polishchuk, M. Salmona, F. Facchiano, K.N. Burger, A. Mironov, A. Luini, and D. Corda. (1999). CtBP/BARS induces fission of Golgi membranes by acylating lysophosphatidic acid. *Nature* 402, 429-433
266. J. Bai and R. E. Pagano. (1997). Measurement of spontaneous transfer and transbilayer movement of BODIPY-labeled lipids in lipid vesicles. *Biochemistry* 36, 8840-8848
267. A. Luna, O.B. Matas, J.A. Martinez-Menarguez, E. Mato, J.M. Duran, J. Ballesta, M. Way and G. Egea. (2002). Regulation of protein transport from the Golgi complex to the endoplasmic reticulum by CDC42 and N-WASP. *Mol.Biol.Cell* 13, 866-879
268. O. B. Matas, J. A. Martinez-Menarguez and G. Egea. (2004). Association of Cdc42/N-WASP/Arp2/3 signaling pathway with Golgi membranes. *Traffic.* 5, 838-846
269. M. D. Brown, B. J. Cornejo, T. B. Kuhn and J. R. Bamberg. (2000). Cdc42 stimulates neurite outgrowth and formation of growth cone filopodia and lamellipodia. *J.Neurobiol.* 43, 352-364
270. M. Corrotte, S. Chasserot-Golaz, P. Huang, G. Du, N.T. Ktistakis, M.A. Frohman, N. Vitale, M.F. Bader and N.J. Grant. (2006). Dynamics and function of phospholipase D and phosphatidic acid during phagocytosis. *Traffic.* 7, 365-377
271. W. E. Balch, B. S. Glick and J. E. Rothman. (1984). Sequential intermediates in the pathway of intercompartmental transport in a cell-free system. *Cell* 39, 525-536
272. J. E. Preiss, R. M. Bell and J. E. Niedel. (1987). Diacylglycerol mass measurements in stimulated HL-60 phagocytes. *J.Immunol.* 138, 1542-1545
273. E. G. BLIGH and W. J. DYER. (1959). A rapid method of total lipid extraction and purification. *Can.J.Biochem.Physiol* 37, 911-917
274. L. Johannes, D. Tenza, C. Antony and B. Goud. (1997). Retrograde transport of KDEL-bearing B-fragment of Shiga toxin. *J.Biol.Chem.* 272, 19554-19561
275. F. Mallard, C. Antony, D. Tenza, J. Salamero, B. Goud and L. Johannes. (1998). Direct pathway from early/recycling endosomes to the Golgi apparatus revealed through the study of shiga toxin B-fragment transport. *J.Cell Biol.* 143, 973-990
276. S.E. Phillips, B. Sha, L. Topalof, Z. Xie, J.G. Alb, V.A. Klenchin, P. Swigart, S. Cockcroft, T.F. Martin, M. Luo and V.A. Bankaitis. (1999). Yeast Sec14p deficient in phosphatidylinositol transfer activity is functional in vivo. *Mol.Cell* 4, 187-197
277. R. P. Huijbregts, L. Topalof and V. A. Bankaitis. (2000). Lipid metabolism and regulation of membrane trafficking. *Traffic.* 1, 195-202
278. A. S. Pappu and G. Hauser. (1983). Propranolol-induced inhibition of rat brain cytoplasmic phosphatidate phosphohydrolase. *Neurochem.Res.* 8, 1565-1575

279. J.E. Bleasdale, N.R. Thakur, R.S. Gremban, G.L. Bundy, F.A. Fitzpatrick, R.J. Smith and S. Bunting. (1990). Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J.Pharmacol.Exp.Ther.* 255, 756-768
280. A. H. Merrill, Jr., M. C. Sullards, E. Wang, K. A. Voss and R. T. Riley. (2001). Sphingolipid metabolism: roles in signal transduction and disruption by fumonisins. *Environ.Health Perspect.* 109 Suppl 2, 283-289
281. E. Wang, W. P. Norred, C. W. Bacon, R. T. Riley and A. H. Merrill, Jr. (1991). Inhibition of sphingolipid biosynthesis by fumonisins. Implications for diseases associated with *Fusarium moniliforme*. *J.Biol.Chem.* 266, 14486-14490
282. W.I. Wu, V.M. McDonough, J.T.Jr. Nickels, J. Ko, A.S. Fischl, T.R. Vales, A.H.Jr. Merrill and G.M. Carman. (1995). Regulation of lipid biosynthesis in *Saccharomyces cerevisiae* by fumonisin B1. *J.Biol.Chem.* 270, 13171-13178
283. S. Schutze, K. Potthoff, T. Machleidt, D. Berkovic, K. Wiegmann and M. Kronke. (1992). TNF activates NF-kappa B by phosphatidylcholine-specific phospholipase C-induced "acidic" sphingomyelin breakdown. *Cell* 71, 765-776
284. G. Egea, F. Lazaro-Dieguez and M. Vilella. (2006). Actin dynamics at the Golgi complex in mammalian cells. *Curr.Opin.Cell Biol.* 18, 168-178
285. E. E. Vaughan and D. A. Dean. (2006). Intracellular trafficking of plasmids during transfection is mediated by microtubules. *Mol.Ther.* 13, 422-428
286. A. Murshid and J. F. Presley. (2004). ER-to-Golgi transport and cytoskeletal interactions in animal cells. *Cell Mol.Life Sci.* 61, 133-145
287. M. J. Lewis and H. R. Pelham. (1992). Ligand-induced redistribution of a human KDEL receptor from the Golgi complex to the endoplasmic reticulum. *Cell* 68, 353-364
288. R. D. Klausner, J. G. Donaldson and J. Lippincott-Schwartz. (1992). Brefeldin A: insights into the control of membrane traffic and organelle structure. *J.Cell Biol.* 116, 1071-1080
289. N. Sciaky, J. Presley, C. Smith, K.J. Zaal, N. Cole, J.E. Moreira, M. Terasaki, E. Siggia and J. Lippincott-Schwartz. (1997). Golgi tubule traffic and the effects of brefeldin A visualized in living cells. *J.Cell Biol.* 139, 1137-1155
290. J. L. Zugaza, J. Sinnett-Smith, J. Van Lint and E. Rozengurt. (1996). Protein kinase D (PKD) activation in intact cells through a protein kinase C-dependent signal transduction pathway. *EMBO J.* 15, 6220-6230
291. R. J. Davis, B. R. Ganong, R. M. Bell and M. P. (1985). sn-1,2-Dioctanoylglycerol. A cell-permeable diacylglycerol that mimics phorbol diester action on the epidermal growth factor receptor and mitogenesis. *J.Biol.Chem.* 260, 1562-1566
292. E. G. Lapetina, B. Reep, B. R. Ganong, R. M. Bell. (1985). Exogenous sn-1,2-diacylglycerols containing saturated fatty acids function as bioregulators of protein kinase C in human platelets. *J.Biol.Chem.* 260, 1358-1361
293. A. M. Vergnano, R. Schlichter and P. Poisbeau. (2007). PKC activation sets an upper limit to the functional plasticity of GABAergic transmission induced by endogenous neurosteroids. *Eur.J.Neurosci.* 26, 1173-1182
294. T. P. Woodsome, A. Polzin, K. Kitazawa, M. Eto and T. Kitazawa. (2006). Agonist- and depolarization-induced signals for myosin light chain phosphorylation and force generation of cultured vascular smooth muscle cells. *J.Cell Sci.* 119, 1769-1780

295. F. Colon-Gonzalez and M. G. Kazanietz. (2006). C1 domains exposed: from diacylglycerol binding to protein-protein interactions. *Biochim.Biophys.Acta* 1761, 827-837
296. J.S. Yang, S.Y. Lee, M. Gao, S. Bourgoïn, P.A. Randazzo, R.T. Premont, V.W. Hsu. (2002). ARFGAP1 promotes the formation of COPI vesicles, suggesting function as a component of the coat. *J.Cell Biol.* 159, 69-78
297. D. Escalante-Alcalde, L. Hernandez, H. Le Stunff, R. Maeda, H.S. Lee, Jr. Gang Cheng, V.A. Sciorra, I. Daar, S. Spiegel, A.J. Morris and C.L. Stewart. (2003). The lipid phosphatase LPP3 regulates extra-embryonic vasculogenesis and axis patterning. *Development* 130, 4623-4637
298. Sundberg,C.; Thodeti,C.K.; Kveiborg,M.; Larsson,C.; Parker,P.; Albrechtsen,R.; Wewer,U.M.. (2004). Regulation of ADAM12 cell-surface expression by protein kinase C epsilon. *J.Biol.Chem.* 279, 51601-51611
299. P. de Figueiredo, R. S. Polizotto, D. Drecktrah and W. J. Brown. (1999). Membrane tubule-mediated reassembly and maintenance of the Golgi complex is disrupted by phospholipase A2 antagonists. *Mol.Biol.Cell* 10, 1763-1782
300. K. E. Morris, L. M. Schang and D. N. Brindley. (2006). Lipid phosphate phosphatase-2 activity regulates S-phase entry of the cell cycle in Rat2 fibroblasts. *J.Biol.Chem.* 281, 9297-9306
301. J. A. Szule, N. L. Fuller and R. P. Rand. (2002). The effects of acyl chain length and saturation of diacylglycerols and phosphatidylcholines on membrane monolayer curvature. *Biophys.J.* 83, 977-984
302. R. Prekeris, R. M. Hernandez, M. W. Mayhew, M. K. White and D. M. Terrian. (1998). Molecular analysis of the interactions between protein kinase C-epsilon and filamentous actin. *J.Biol.Chem.* 273, 26790-26798
303. M.C. Souroujon, L. Yao, H. Chen, G. Endemann, H. Khaner, V. Geeraert, D. Schechtman, A.S. Gordon, I. Diamond and D. Mochly-Rosen. (2004). State-specific monoclonal antibodies identify an intermediate state in epsilon protein kinase C activation.*J.Biol.Chem.* 279, 17617-17624